

## PCT COOPERATION TREATY

## PCT

## NOTIFICATION RELATING TO PRIORITY CLAIM

(PCT Rules 26bis.1 and 26bis.2 and  
Administrative Instructions, Sections 402 and 409)

From the INTERNATIONAL BUREAU

To:

FITZNER, Uwe  
Lintorfer Str. 10  
D-40878 Ratingen  
ALLEMAGNE

Date of mailing (day/month/year) 13 July 2000 (13.07.00)	<b>IMPORTANT NOTIFICATION</b>
Applicant's or agent's file reference 3377/99	
International application No. PCT/EP00/02701	International filing date (day/month/year) 28 March 2000 (28.03.00)
Applicant BASF PLANT SCIENCE GMBH et al	

The applicant is hereby notified of the following in respect of the priority claim(s) made in the international application.

1. ☒ **Correction of priority claim.** In accordance with the applicant's notice received on: 22 June 2000 (22.06.00), the following priority claim has been corrected to read as follows:

US 07 February 2000 (07.02.00) 60/180,687

- ☐ even though the indication of the number of the earlier application is missing.  
☐ even though the following indication in the priority claim is not the same as the corresponding indication appearing in the priority document:

2. ☐ **Addition of priority claim.** In accordance with the applicant's notice received on: , the following priority claim has been added:

- ☐ even though the indication of the number of the earlier application is missing.  
☐ even though the following indication in the priority claim is not the same as the corresponding indication appearing in the priority document:

3. ☐ As a result of the correction and/or addition of (a) priority claim(s) under items 1 and/or 2, the (earliest) priority date is:

4. ☐ **Priority claim considered not to have been made.**

- ☐ The applicant failed to respond to the Invitation under Rule 26bis.2(a) (Form PCT/IB/316) within the prescribed time limit.  
☐ The applicant's notice was received after the expiration of the prescribed time limit under Rule 26bis.1(a).  
☐ The applicant's notice failed to correct the priority claim so as to comply with the requirements of Rule 4.10.

The applicant may, before the technical preparations for international publication have been completed and subject to the payment of a fee, request the International Bureau to publish, together with the international application, information concerning the priority claim. See Rule 26bis.2(c) and the PCT Applicant's Guide, Volume I, Annex B2(1B).

5. ☒ In case where multiple priorities have been claimed, the above item(s) relate to the following priority claim(s):  
 US 07 February 2000 (07.02.00) 60/180,687

6. A copy of this notification has been sent to the receiving Office and

- ☒ to the International Searching Authority (where the international search report has not yet been issued).  
☒ the designated Offices (which have already been notified of the receipt of the record copy).

The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland  Facsimile No. (41-22) 740.14.35	Authorized officer  Aino Metcalfe  Telephone No. (41-22) 338.83.38
--	--

## PATENT COOPERATION TREATY

PCT

## NOTIFICATION OF ELECTION

(PCT Rule 61.2)

From the INTERNATIONAL BUREAU

To:

Commissioner  
US Department of Commerce  
United States Patent and Trademark  
Office, PCT  
2011 South Clark Place Room  
CP2/5C24  
Arlington, VA 22202  
ETATS-UNIS D'AMERIQUE

in its capacity as elected Office

Date of mailing (day/month/year) 21 February 2001 (21.02.01)	
International application No. PCT/EP00/02701	Applicant's or agent's file reference 3377/99
International filing date (day/month/year) 28 March 2000 (28.03.00)	Priority date (day/month/year) 01 April 1999 (01.04.99)
Applicant DAHLQVIST, Anders et al	

1. The designated Office is hereby notified of its election made:

☒ in the demand filed with the International Preliminary Examining Authority on:

11 October 2000 (11.10.00)

☐ in a notice effecting later election filed with the International Bureau on:2. The election ☒ was☐ was not

made before the expiration of 19 months from the priority date or, where Rule 32 applies, within the time limit under Rule 32.2(b).

The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland	Authorized officer C. Cupello
Facsimile No.: (41-22) 740.14.35	Telephone No.: (41-22) 338.83.38

# PATENT COOPERATION TREATY

## PCT

REC'D 08 JUN 2001	
WIPO	PCT

### INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)

*14*

Applicant's or agent's file reference <b>3377/99 PCT</b>	<b>FOR FURTHER ACTION</b>		See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)
International application No. <b>PCT/EP00/02701</b>	International filing date (day/month/year) <b>28/03/2000</b>	Priority date (day/month/year) <b>01/04/1999</b>	
International Patent Classification (IPC) or national classification and IPC <b>C12N15/54</b>			
Applicant <b>BASF PLANT SCIENCE GmbH et al.</b>			

1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.



2. This REPORT consists of a total of 12 sheets, including this cover sheet.

- ☐ This report is also accompanied by ANNEXES, i.e. sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).

These annexes consist of a total of sheets.

3. This report contains indications relating to the following items:

- I ☒ Basis of the report
- II ☒ Priority
- III ☒ Non-establishment of opinion with regard to novelty, inventive step and industrial applicability
- IV ☒ Lack of unity of invention
- V ☒ Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
- VI ☒ Certain documents cited
- VII ☒ Certain defects in the international application
- VIII ☒ Certain observations on the international application

Date of submission of the demand  <b>11/10/2000</b>	Date of completion of this report  <b>06.06.2001</b>
Name and mailing address of the international preliminary examining authority:  <b>European Patent Office</b> <b>D-80298 Munich</b> <b>Tel. +49 89 2399 - 0 Tx: 523656 epmu d</b> <b>Fax: +49 89 2399 - 4465</b>	Authorized officer  Page, M  Telephone No. +49 89 2399 7322 <div style="text-align: right;">  </div>

# INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/EP00/02701

## I. Basis of the report

1. With regard to the **elements** of the international application (*Replacement sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to this report since they do not contain amendments (Rules 70.16 and 70.17)*):

### Description, pages:

1-32 as originally filed

### Claims, No.:

1-22 as originally filed

### Drawings, sheets:

1/6-6/6 as originally filed

### Sequence listing part of the description, pages:

1-45 (SEQ ID NOs. 1-15), as originally filed

2. With regard to the **language**, all the elements marked above were available or furnished to this Authority in the language in which the international application was filed, unless otherwise indicated under this item.

These elements were available or furnished to this Authority in the following language: , which is:

- ☐ the language of a translation furnished for the purposes of the international search (under Rule 23.1(b)).
- ☐ the language of publication of the international application (under Rule 48.3(b)).
- ☐ the language of a translation furnished for the purposes of international preliminary examination (under Rule 55.2 and/or 55.3).

3. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international preliminary examination was carried out on the basis of the sequence listing:

- ☒ contained in the international application in written form.
- ☐ filed together with the international application in computer readable form.
- ☐ furnished subsequently to this Authority in written form.
- ☒ furnished subsequently to this Authority in computer readable form.
- ☒ The statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.
- ☒ The statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.

4. The amendments have resulted in the cancellation of:

# INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/EP00/02701

- ☐ the description, pages:
- ☐ the claims, Nos.:
- ☐ the drawings, sheets:

5. ☐ This report has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed (Rule 70.2(c)):

*(Any replacement sheet containing such amendments must be referred to under item 1 and annexed to this report.)*

6. Additional observations, if necessary:

### III. Non-establishment of opinion with regard to novelty, inventive step and industrial applicability

1. The questions whether the claimed invention appears to be novel, to involve an inventive step (to be non-obvious), or to be industrially applicable have not been examined in respect of:

- ☐ the entire international application.
- ☒ claims Nos. 22 (partially).

because:

- ☐ the said international application, or the said claims Nos. relate to the following subject matter which does not require an international preliminary examination (*specify*):
- ☐ the description, claims or drawings (*indicate particular elements below*) or said claims Nos. are so unclear that no meaningful opinion could be formed (*specify*):
- ☒ the claims, or said claims Nos. 22 (partially) are so inadequately supported by the description that no meaningful opinion could be formed.
- ☐ no international search report has been established for the said claims Nos. .

2. A meaningful international preliminary examination cannot be carried out due to the failure of the nucleotide and/or amino acid sequence listing to comply with the standard provided for in Annex C of the Administrative Instructions:

- ☐ the written form has not been furnished or does not comply with the standard.
- ☐ the computer readable form has not been furnished or does not comply with the standard.

### IV. Lack of unity of invention

1. In response to the invitation to restrict or pay additional fees the applicant has:

# INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/EP00/02701

- ☐ restricted the claims.
  - ☐ paid additional fees.
  - ☐ paid additional fees under protest.
  - ☐ neither restricted nor paid additional fees.
2. ☒ This Authority found that the requirement of unity of invention is not complied and chose, according to Rule 68.1, not to invite the applicant to restrict or pay additional fees.
3. This Authority considers that the requirement of unity of invention in accordance with Rules 13.1, 13.2 and 13.3 is
- ☐ complied with.
  - ☒ not complied with for the following reasons:  
**see separate sheet**
4. Consequently, the following parts of the international application were the subject of international preliminary examination in establishing this report:
- ☒ all parts.
  - ☐ the parts relating to claims Nos. .

## V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

### 1. Statement

Novelty (N)	Yes:	Claims	9-19, 21, 22 (all partially)
	No:	Claims	1-19, 21, 22 (all partially)
Inventive step (IS)	Yes:	Claims	9-19, 21, 22 (all partially)
	No:	Claims	1-19, 21, 22 (all partially)
Industrial applicability (IA)	Yes:	Claims	1-22
	No:	Claims	

### 2. Citations and explanations **see separate sheet**

## VI. Certain documents cited

### 1. Certain published documents (Rule 70.10)

and / or

### 2. Non-written disclosures (Rule 70.9)

**INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT**

International application No. PCT/EP00/02701

---

**see separate sheet**

**VII. Certain defects in the international application**

The following defects in the form or contents of the international application have been noted:  
**see separate sheet**

**VIII. Certain observations on the international application**

The following observations on the clarity of the claims, description, and drawings or on the question whether the claims are fully supported by the description, are made:  
**see separate sheet**

**INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT - SEPARATE SHEET**

---

International application No. PCT/EP00/02701

The application concerns the provision of a yeast and plant polypeptide and polynucleotide sequences allegedly corresponding to diacylglycerol acyltransferases. Function is shown for *Saccharomyces cerevisiae* sequences, but neither the function nor any structural relationship to the *Saccharomyces* sequences making such a function plausible are demonstrated for the other full-length and partial sequences.

**Re Item II**

**Priority**

After considering the priority document, the documents cited "P, X" in the search report are not considered relevant for the examination of novelty and inventive step.

**Re Item III**

**Non-establishment of opinion with regard to novelty, inventive step and industrial applicability**

**Claim 18 (claim 22 as originally filed)** seeks protection for cells or organisms with altered PDAT activity, "wherein the altered PDAT activity is characterized by an alteration in gene expression, catalytic activity and/or regulation of activity of the enzyme". No reference could be found in the description for alterations to the catalytic activity or regulation of PDAT activity and claim 18 (partially) is therefore considered to lack meaningful support from the description. The claim has only been examined with respect to alterations in gene expression.

**Re Item IV**

**Lack of Unity of Invention**

An international application must relate to one invention only or to a group of inventions so linked as to form a single general inventive concept. Unity of invention is fulfilled only when there is a technical relationship between the inventions involving one or more of the same or corresponding special technical features. Special technical features are such features that define the contribution of the claimed



invention over the prior art.

The identified 8 inventions relate to a group of sequences with the claimed technical feature of being diacylglycerol acyltransferases as the sole common link. However, this feature cannot be considered to constitute a special technical feature because it does not define a contribution over the prior art: SEQ ID NOs. 2, 3, 9, 16, 20 and 22 have all been previously disclosed in their entirety (D1, D2 and D3).

Although the prior art does not disclose the function of the encoded enzymes, they do disclose the nucleic acid and polypeptide sequences of the respective claimed SEQ ID NOs. The encoded enzyme will have the activity claimed in claim 1, regardless of whether or not this is disclosed in the prior art.

The application therefore does not meet the requirements of Rule 13.2 PCT in that there is no common special technical feature linking the 8 inventions of the application, these being:

**Invention I      Claims 5, 6, 8-22 (all partially) and 1-3 (completely) (formerly claims 1, 3, 6, 7, 9, 11-27 (all partially) 2 and 4 (completely))**

Enzymes catalysing the acyl-CoA-independent transfer of fatty acids to diacylglycerol in the production of triacylglycerol from *Saccharomyces cerevisiae* and corresponding to polypeptides with SEQ ID NOs. 2, 16, 20 and 22, encoded by polynucleotides SEQ ID NOs. 1, 19 and 21, fragments, derivatives, alleles, homologs and isoenzymes, the corresponding polynucleotide sequences, portions, derivatives, alleles and homologs of the polynucleotide sequence, expression vectors, transgenic cells and organisms, processes for the production of triacylglycerol using such cells/organisms, the product of such a process and the use of the enzymes and polynucleotides in such processes.

**Invention II      Claims 4-6 and 8-22 (all partially) (formerly claims 1, 3, 5-9 and 11-27 (all partially))**

As invention I with SEQ ID NOs. 3, 13 and 23 from *Schizosaccharomyces pombe*.

**Invention III      Claims 4-22 (all partially) (formerly claims 1, 3 and 5-27 (all partially))**

As invention I with SEQ ID NOs. 4-6, 18, 24, 25 from *Arabidopsis thaliana*.

**Invention IV**    **Claims 4, 5 and 7-22 (all partially) (formerly claims 1, 3 and 5-27 (all partially))**

As invention I with SEQ ID NOs. 7, 8, 26 and 27 from *Zea mays*.

**Invention V**    **Claims 5 and 7-22 (all partially) (formerly claims 1, 3, 6-8 and 10-27 (all partially))**

As invention I with SEQ ID NOs. 9 and 28 from *Neurospora crassa*.

**Invention VI**    **Claims 4-6 and 8-22 (all partially) (formerly claims 1, 3, 5-9 and 11-27 (all partially))**

As invention I with SEQ ID NOs. 10, 14, 17 and 29 from *Arabidopsis thaliana*.

**Invention VII**    **Claims 4-6 and 8-22 (all partially) (formerly claims 1, 3, 5-9 and 11-27 (all partially))**

As invention I with SEQ ID NOs. 11, 15 and 30 from *Arabidopsis thaliana*.

**Invention VIII**    **Claims 5 and 7-22 (all partially) (formerly claims 1, 3 and 5-27 (all partially))**

As invention I with SEQ ID NOs. 12 and 31 from *Lycopersicon esculentum*.

**Re Item V**

**Reasoned statement under Rule 66.2(a)(ii) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement**

**1) Reference is made to the following documents:**

D1: PETER VERHASSELT ET AL.: 'Twelve open reading frames revealed in the 23.6kb segment flanking the centromere on the *Saccharomyces cerevisiae* chromosome XIV right arm' YEAST, vol. 10, no. 7, July 1994 (1994-07), pages 1355-1361, XP002112572 -& Swissprot Database Entry Yn84\_Yeast Accession number P40345; 1 February 1995 XP002112574

D2: DATABASE EMBL [Online] Database Entry SPBC776, 21 January 1999 (1999-01-21) LYNE M. ET AL.: 'S. pombe chromosome II cosmid c776' Database accession no. AL035263 XP002150203

D3: DATABASE EMBL [Online] Database Entry AI398644, 10 February 1999

(1999-02-10) XP002150204 & MARY ANNE NELSON ET AL.: 'Expressed sequences from conidial, mycelial, and sexual stages of *Neurospora crassa*' FUNGAL GENETICS AND BIOLOGY, vol. 21, 1997, pages 348-363, XP000952173

D4: KEITH STOBART ET AL.: 'Triacylglycerols are synthesized and utilized by transacylation reactions in microsomal preparations of developing safflower (*Carthamus tinctorius* L.) seeds' PLANTA, vol. 203, no. 1, 1997, pages 58-66, XP002112573

D5: WO 98 55631 A (CALGENE LLC) 10 December 1998 (1998-12-10)

2) **Novelty - Art.33(1) and (2) PCT:**

**Invention I      Claims 5, 6, 8 (all partially) and 1-3 (completely)**

**Claims 5, 6, 8 (partially), and 1-3 (completely)** lack novelty in light of the sequence with the accession number P40345 provided by D1 (identified therein as N2042) which, according to the description of the present application, encodes an acyl-CoA-independent acyltransferase. Although D1 does not disclose the function of the encoded enzyme, a polynucleotide or polypeptide sequence is not rendered novel by the discovery of its function. The disclosed sequence is 100% identical to SEQ ID NO. 2 over the whole length of the protein.

**Inventions II-VIII      Claims 4-8 (all partially)**

**Claims 4-8 (partially)** lack novelty in light of the sequences provided by D1, D2 and D3 which, according to the description, are polypeptides and polynucleotides corresponding to phospholipid:diacylglycerol acyltransferases. As stated previously, identifying the function of known polypeptides does not render the polypeptides novel.

The description, for example, defines a "functional fragment" on page 4 lines 30-32 as being "any polypeptide sequence which shows specific enzyme activity of a *PDAT*" The enzyme N2042 disclosed in D1 clearly possesses such activity and thus the claims lack novelty.

Similarly, allelic variants are understood to be "any different nucleotide sequence which encodes a polypeptide with a functionally different function" and having an undisclosed number of substitutions, additions or deletions (page 5 lines 28). Again,

the protein of D1 clearly fulfills these requirements.

The definition provided on page 6 lines 17 and 18 for the term "isoenzyme" meets the same objections.

Furthermore, the definition in the description for the term "portion" is meant to include any nucleotide sequence which shows specific activity of a PDAT" (page 5 lines 7-17). The term includes within its scope the polynucleotide sequences A1398644 of D3 for example.

**Inventions I-VIII      Claims 9-22**

**Claims 9-19, 21 and 22 (partially)** appear to be novel in light of the cited prior art. although polynucleotide and polypeptide sequences according to the claimed invention have been disclosed (e.g. D1 sequence N2042, D2 sequence O94680, D3), these documents do not disclose gene constructs, vectors, transgenic cells or the use of such in the production of triacylglycerol.

**Claim 20 (partially)** lacks novelty in light of D4, which discloses triacylglycerol made with an acyl-Co-A independent acyltransferase (D4 page 59 left-hand column paragraph 1). Even if the claim were restricted to triacylglycerol made using novel subject matter, the Applicant would need to show how this product differs from previously disclosed subject matter, as a product is not rendered novel by a new method for making it.

**3) Inventive Step - Art.33(1) and (3) PCT:**

The following comments on inventive step are confined to subject matter which could be acknowledged as being novel, or for which novelty could potentially be restored as outlined supra.

**Invention I      Claims 9-19, 21 and 22 (all partially)**

The closest prior art is document D5, which discloses a the polypeptide and polynucleotide sequences for an acyl-Co-A dependent plant diacylglycerol acyltransferase as well as the use of the sequences in engineering plants with altered triacylglycerol content (D5 page 3 line 22 to page 5 line 20).

In the light of the prior art, the technical problem can be regarded as the provision of further polynucleotide and polypeptide sequences encoding enzymes that can alter the triacylglycerol content of cells or organisms expressing them.

**Claims 9-19, 21 and 22** appear to be inventive in light of the cited prior art, which does not disclose the enzyme activity of SEQ ID NO. 2. There is therefore no motivation to combine the teaching of D5 with that of D1 disclosing the sequence N2042.

**Inventions II-VIII 9-19, 21 and 22 (all partially)**

Again, the closest prior art is document D5, which discloses a the polypeptide and polynucleotide sequences for an acyl-Co-A dependent plant diacylglycerol acyltransferase as well as the use of the sequences in engineering plants with altered triacylglycerol content (D5 page 3 line 22 to page 5 line 20).

In the light of the prior art, the technical problem can be regarded as the provision of further polynucleotide and polypeptide sequences encoding enzymes that can alter the triacylglycerol content of cells or organisms expressing them.

It cannot be seen how inventive step can be recognised for **claims 9-19, 21 and 22**. Although function has been demonstrated for the enzyme encoded by SEQ ID NO. 1, no such function has been demonstrated for the sequences from other species, nor has the Applicant shown that there is a structural relationship between the sequences of Invention I and those of Inventions II-VIII that would make such a function plausible. This is true for the full-length sequences as well as the partial sequences disclosed in the application.

**Re Item VII**

**Certain defects in the international application**

- a) Contrary to the requirements of Rule 5.1(a)(ii) PCT, the relevant background art disclosed in the documents D1-D5 are not mentioned in the description, nor are these documents identified therein.

**INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT - SEPARATE SHEET**

---

International application No. PCT/EP00/02701

**Rule VIII**

**Certain observations on the international application**

- a) Several of the SEQ ID NOs. appear to be identical duplicates of each other, resulting in a lack of conciseness as required by Article 6 PCT. The unnecessary duplicates should be removed.

# PATENT COOPERATION TREATY

# PCT

## INTERNATIONAL SEARCH REPORT

(PCT Article 18 and Rules 43 and 44)

Applicant's or agent's file reference <b>3377/99 PCT</b>	<div style="display: flex; justify-content: space-between;"> <div style="text-align: center;"> <b>FOR FURTHER ACTION</b> </div> <div style="font-size: small;">             see Notification of Transmittal of International Search Report (Form PCT/ISA/220) as well as, where applicable, item 5 below.           </div> </div>	
International application No. <b>PCT/EP 00/ 02701</b>	International filing date (day/month/year) <div style="text-align: center;"><b>23/03/2000</b></div>	(Earliest) Priority Date (day/month/year) <div style="text-align: center;"><b>01/04/1999</b></div>
Applicant  <b>BASF PLANT SCIENCE GmbH</b>		

This International Search Report has been prepared by this International Searching Authority and is transmitted to the applicant according to Article 18. A copy is being transmitted to the International Bureau.

This International Search Report consists of a total of 3 sheets.

☒ It is also accompanied by a copy of each prior art document cited in this report.

### 1. Basis of the report

a. With regard to the **language**, the international search was carried out on the basis of the international application in the language in which it was filed, unless otherwise indicated under this item.

☐ the international search was carried out on the basis of a translation of the international application furnished to this Authority (Rule 23.1(b)).

b. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international search was carried out on the basis of the sequence listing :

☒ contained in the international application in written form.

☐ filed together with the international application in computer readable form.

☐ furnished subsequently to this Authority in written form.

☒ furnished subsequently to this Authority in computer readable form.

☒ the statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.

☒ the statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished

2. ☐ **Certain claims were found unsearchable** (See Box I).

3. ☐ **Unity of Invention is lacking** (see Box II).

4. With regard to the **title**,

☐ the text is approved as submitted by the applicant.

☒ the text has been established by this Authority to read as follows:

**ENZYMES OF THE BIOSYNTHETIC PATHWAY FOR THE PRODUCTION OF TRIACYLGLYCEROL AND RECOMBINANT DNA MOLECULES ENCODING THESE ENZYMES**

5. With regard to the **abstract**,

☒ the text is approved as submitted by the applicant.

☐ the text has been established, according to Rule 38.2(b), by this Authority as it appears in Box III. The applicant may, within one month from the date of mailing of this international search report, submit comments to this Authority.

6. The figure of the **drawings** to be published with the abstract is Figure No.

☒ as suggested by the applicant.

☐ because the applicant failed to suggest a figure.

☐ because this figure better characterizes the invention.

1  
☐ None of the figures.

## INTERNATIONAL SEARCH REPORT

International Application No

T/EP 00/02701

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/54 C12N9/10 C12N15/81 C12N15/82 C12N1/16  
C12N5/10 A01K67/027 C12P7/64

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N A01K C12P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

STRAND, EPO-Internal, WPI Data, MEDLINE, CHEM ABS Data, BIOSIS, EMBL

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	PETER VERHASSELT ET AL.: "Twelve open reading frames revealed in the 23.6kb segment flanking the centromere on the <i>Saccharomyces cerevisiae</i> chromosome XIV right arm" YEAST, vol. 10, no. 7, July 1994 (1994-07), pages 1355-1361, XP002112572 abstract; table 2	1-23,27
X	-& Swissprot Database Entry Yn84_Yeast Accession number P40345; 1 February 1995 XP002112574 the whole document --- -/--	1-23,27



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

## ° Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&amp;" document member of the same patent family

Date of the actual completion of the international search

17 October 2000

Date of mailing of the international search report

30/10/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Montero Lopez, B



## INTERNATIONAL SEARCH REPORT

International Application No

T/EP 00/02701

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE EMBL 'Online!  Database Entry SPBC776,  21 January 1999 (1999-01-21)  LYNE M. ET AL.: "S. pombe chromosome II  cosmid c776"  Database accession no. AL035263  XP002150203  the whole document</p>	1-23,27
X	<p>DATABASE EMBL 'Online!  Database Entry AI398644,  10 February 1999 (1999-02-10)  XP002150204  the whole document  &amp; MARY ANNE NELSON ET AL.: "Expressed  sequences from conidial, mycelial, and  sexual stages of Neurospora crassa "  FUNGAL GENETICS AND BIOLOGY,  vol. 21, 1997, pages 348-363, XP000952173</p>	1-23,27
X	<p>KEITH STOBART ET AL.: "Triacylglycerols  are synthesized and utilized by  transacylation reactions in microsomal  preparations of developing safflower  (Carthamus tinctorius L.) seeds"  PLANTA,  vol. 203, no. 1, 1997, pages 58-66,  XP002112573  page 58, right-hand column, last paragraph  -page 59, left-hand column, paragraph 1  page 63, right-hand column, paragraph 2</p>	25
A	<p>WO 98 55631 A (CALGENE LLC)  10 December 1998 (1998-12-10)  page 9, line 36 -page 10, line 7  page 12, line 28 -page 13, line 18  page 14, line 34 -page 15, line 13  page 20, line 5 -page 25, line 4</p>	1-27
P,X	<p>DATABASE SWALL 'Online!  Database Entry 094680,  1 May 1999 (1999-05-01)  LYNE M. ET AL.: "hypothetical 69.7 kDa  protein C776.14 in chromosome II"  Database accession no. 094680  XP002150205  the whole document</p>	1-23,27

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

EP 00/02701


Patent document cited in search report	Publication date	Patent family member(s)	Publication date
W0 9855631 A	10-12-1998	CN 1266460 T EP 1003882 A	13-09-2000 31-05-2000



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : <b>C12N 15/54, 9/10, 15/81, 15/82, 1/16, 5/10, A01N 27/067, C12P 7/64</b>		<b>A2</b>	(11) International Publication Number: <b>WO 00/60095</b>
			(43) International Publication Date: 12 October 2000 (12.10.00)
(21) International Application Number: PCT/EP00/02701		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 28 March 2000 (28.03.00)			
(30) Priority Data: 99106656.4 1 April 1999 (01.04.99) EP 99111321.8 10 June 1999 (10.06.99) EP 60/180,687 7 February 2000 (07.02.00) US			
(71) Applicant (for all designated States except US): BASF PLANT SCIENCE GMBH [DE/DE]; D-67056 Ludwigshafen (DE).			
(72) Inventors; and		Published	
(75) Inventors/Applicants (for US only): DAHLQVIST, Anders [SE/SE]; Hemmansvägen 2, S-244 66 Furulund (SE). STAHL, Ulf [SE/SE]; Liljegatan 7b, S-753 24 Uppsala (SE). LENMAN, Marit [SE/SE]; Revingegatan 13a, S-223 59 Lund (SE). BANAS, Antoni [SE/PL]; Wiolinowa 14, PL-08110 Siedlce (PL). RONNE, Hans [SE/SE]; Dirigentvägen 169, S-756 54 Uppsala (SE). STYMNE, Sten [SE/SE]; Torrlösa 1380, S-269 90 Svalöv (SE).		Without international search report and to be republished upon receipt of that report.	
(74) Agent: FITZNER, Uwe; Lintorfer Str. 10, D-40878 Ratingen (DE).			

(54) Title: A NEW CLASS OF ENZYMES IN THE BIOSYNTHETIC PATHWAY FOR THE PRODUCTION OF TRIACYLGLYCEROL AND RECOMBINANT DNA MOLECULES ENCODING THESE ENZYMES



## (57) Abstract

The present invention relates to the isolation, identification and characterization of nucleotide sequences encoding an enzyme catalysing the transfer of fatty acids from phospholipids to diacylglycerol in the biosynthetic pathway for the production of triacylglycerol, to the said enzymes and a process for the production of triacylglycerols.

*FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

A NEW CLASS OF ENZYMES IN THE BIOSYNTHETIC PATHWAY FOR THE  
PRODUCTION OF TRIACYLGLYCEROL AND RECOMBINANT DNA  
MOLECULES ENCODING THESE ENZYMES

- 5 The present invention relates to the isolation, identification and characterization of recombinant DNA molecules encoding enzymes catalysing the transfer of fatty acids from phospholipids to diacylglycerol in the biosynthetic pathway for the production of triacylglycerol.
- 10 Triacylglycerol (TAG) is the most common lipid-based energy reserve in nature. The main pathway for synthesis of TAG is believed to involve three sequential acyl-transfers from acyl-CoA to a glycerol backbone (1, 2). For many years, acyl-CoA : diacylglycerol acyltransferase (DAGAT), which catalyzes the third acyl transfer reaction, was thought to be the only unique enzyme involved in
- 15 TAG synthesis. It acts by diverting diacylglycerol (DAG) from membrane lipid synthesis into TAG (2). Genes encoding this enzyme were recently identified both in the mouse (3) and in plants (4, 5), and the encoded proteins were shown to be homologous to acyl-CoA : cholesterol acyltransferase (ACAT). It was also recently reported that another DAGAT exists in the oleaginous fungus
- 20 *Mortierella ramanniana*, which is unrelated to the mouse DAGAT, the ACAT gene family or to any other known gene (6).

The instant invention relates to novel type of enzymes and their encoding genes for transformation. More specifically, the invention relates to use of a

25 type of genes encoding a not previously described type of enzymes hereinafter designated phospholipid:diacylglycerol acyltransferases (PDAT), whereby this enzyme catalyses an acyl-CoA-independent reaction. The said type of genes expressed alone in transgenic organisms will enhance the total amount of oil (triacylglycerols) produced in the cells. The PDAT genes, in combination with a

30 gene for the synthesis of an uncommon fatty acid will, when expressed in transgenic organisms, enhance the levels of the uncommon fatty acids in the triacylglycerols.

There is considerable interest world-wide in producing chemical feedstock, such as fatty acids, for industrial use from renewable plant resources rather than non-renewable petrochemicals. This concept has broad appeal to  
5 manufacturers and consumers on the basis of resource conservation and provides significant opportunity to develop new industrial crops for agriculture.

There is a diverse array of unusual fatty acids in oils from wild plant species and these have been well characterised. Many of these acids have industrial  
10 potential and this has led to interest in domesticating relevant plant species to enable agricultural production of particular fatty acids.

Development in genetic engineering technologies combined with greater understanding of the biosynthesis of unusual fatty acids now makes it possible  
15 to transfer genes coding for key enzymes involved in the synthesis of a particular fatty acid from a wild species into domesticated oilseed crops. In this way individual fatty acids can be produced in high purity and quantities at moderate costs.

20 In all crops like rape, sunflower, oilpalm etc., the oil (i.e. triacylglycerols) is the most valuable product of the seeds or fruits and other compounds like starch, protein, and fibre is regarded as by-products with less value. Enhancing the quantity of oil per weight basis at the expense of other compounds in oil crops would therefore increase the value of crop. If genes, regulating the allocation of  
25 reduced carbon into the production of oil can be up-regulated, the cells will accumulate more oil on the expense of other products. Such genes might not only be used in already high oil producing cells, such as oil crops, but could also induce significant oil production in moderate or low oil containing crops such as e.g. soy, oat, maize, potato, sugarbeets, and turnips as well as in  
30 micro-organisms.

## Summary of the invention

Many of the unusual fatty acids of interest, e.g. medium chain fatty acids, hydroxy fatty acids, epoxy fatty acids and acetylenic fatty acids, have physical properties that are distinctly different from the common plant fatty acids. The present inventors have found that, in plant species naturally accumulating these uncommon fatty acids in their seed oil (i.e. triacylglycerol), these acids are absent, or present in very low amounts in the membrane (phospho)lipids of the seed. The low concentration of these acids in the membrane lipids is most likely a prerequisite for proper membrane function and thereby for proper cell functions. One aspect of the invention is that seeds of transgenic crops can be made to accumulate high amounts of uncommon fatty acids if these fatty acids are efficiently removed from the membrane lipids and channelled into seed triacylglycerols.

The inventors have identified a novel class of enzymes in plants catalysing the transfer of fatty acids from phospholipids to diacylglycerol in the production of triacylglycerol through an acyl-CoA-independent reaction and that these enzymes (phospholipid:diacylglycerol acyltransferases, abbreviated as PDAT) are involved in the removal of hydroxylated, epoxygenated fatty acids, and probably also other uncommon fatty acids such as medium chain fatty acids, from phospholipids in plants.

This enzyme reaction was shown to be present in microsomal preparations from baker's yeast (*Saccharomyces cerevisiae*). The instant invention further pertains to an enzyme comprising an amino acid sequence as set forth in SEQ ID No. 2 or a functional fragment, derivate, allele, homolog or isoenzyme thereof. A so called 'knock out' yeast mutant, disrupted in the respective gene was obtained and microsomal membranes from the mutant was shown to totally lack PDAT activity. Thus, it was proved that the disrupted gene encodes a PDAT enzyme (SEQ ID NO. 1 and 2). Furtherm, this PDAT enzyme is

characterized through the amino acid sequence as set forth in SEQ ID NO 2 containing a lipase motif of the conserved sequence string FXKWVEA.

The instant invention pertains further to an enzyme comprising an amino acid sequence as set forth in SEQ ID NO. 1a, 2b or 5a or a functional fragment, 5 derivate, allele, homolog or isoenzyme thereof.

Further genes and/or proteins of so far unknown function were identified and are contemplated within the scope of the instant invention. A gene from *Schizosaccharomyces pombe*, SPBC776.14 (SEQ ID NO. 3), a putative open reading frame CAA22887 of the SPBC776.14 (SEQ ID NO. 13) were identified.

10 Further *Arabidopsis thaliana* genomic sequences (SEQ ID NO. 4, 10 and 11) coding for putative proteins were identified, as well as a putative open reading frame AAC80628 from the *A. thaliana* locus AC 004557 (SEQ ID NO. 14) and a putative open reading frame AAD10668 from the *A. thaliana* locus AC 003027 (SEQ ID NO. 15) were identified.

15 Also, a partially sequenced cDNA clone from *Neurospora crassa* (SEQ ID NO. 9) and a *Zea mays* EST (Extended Sequence Tac) clone (SEQ ID NO. 7) and corresponding putative amino acid sequence (SEQ ID NO. 8) were identified. Finally, two cDNA clones were identified, one *Arabidopsis thaliana* EST (SEQ ID NO. 5 and corresponding predicted amino acid sequence SEQ ID NO. 6) 20 and a *Lycopersicon esculentum* EST clone (SEQ ID NO. 12) were identified.

Further, enzymes designated as PDAT comprising an amino acid sequence selected from the group consisting of sequences as set forth in SEQ ID NO 2a, 3a, 5b, 6 or 7b containing a lipase motif FXKWVEA are contemplated within the scope of the invention. Moreover, an enzyme comprising an amino acid 25 sequence encoded through a nucleotide sequence, a portion, derivate, allele or homolog thereof selected from the group consisting of sequences as set forth in SEQ ID No. 1, 1b, 3, 3b, 4, 4a, 4b, 5, 5b, 6b, 7, 8b, 9, 9b, 10, 10b, 11, 11b or 12 or a functional fragment, derivate, allele, homolog or isoenzyme of the enzyme encoding amino acid sequence are included within the scope of the 30 invention.



A functional fragment of the instant enzyme is understood to be any polypeptide sequence which shows specific enzyme activity of a phospholipid:diacylglycerol acyltransferase (PDAT). The length of the functional fragment can for example vary in a range from about  $660 \pm 10$  amino acids to  $660 \pm 250$  amino acids, preferably from about  $660 \pm 50$  to  $660 \pm 100$  amino acids, whereby the „basic number“ of 660 amino acids corresponds in this case to the polypeptide chain of the PDAT enzyme of SEQ ID NO. 2 encoded by a nucleotide sequence according to SEQ ID NO. 1. Consequently, the „basic number“ of functional fulllength enzyme can vary in correspondance to the encoding nucleotide sequence.

A portion of the instant nucleotide sequence is meant to be any nucleotide sequence encoding a polypeptid which shows specific activity of a phospholipid:diacylglycerol acyltransferase (PDAT). The length of the nucleotide portion can vary in a wide range of about several hundreds of nucleotides based upon the coding region of the gene or a highly conserved sequence. For example the length varies in a range form about  $1900 \pm 10$  to  $1900 \pm 1000$  nucleotides, preferably form about  $1900 \pm 50$  to  $1900 \pm 700$  and more preferably form about  $1900 \pm 100$  to  $1900 \pm 500$  nucleotides. whereby the „basic number“ of 1900 nucleotides corresponds in this case to the encoding nucleotide sequence of the PDAT enzyme of SEQ ID NO. 1. Consequently, the „basic number“ of functional fulllength gene can vary.

An allelic variant of the instant nucleotide sequence is understood to be any different nucleotide sequence which encodes a polypeptide with a functionally equivalent function. The alleles pertain naturally occuring variants of the instant nucleotide sequences as well as synthetic nucleotide sequences produced by methods known in the art. Contemplated are even altered nucleotide sequences which result in an enzyme with altered activity and/or regulation or which is resistant against specific inhibitors. The instant invention further includes natural or synthetic mutations of the originally isolated nucleotide

sequences. These mutations can be substitution, addition, deletion, inversion or insertion of one or more nucleotides.

A homologous nucleotide sequence is understood to be a complementary  
5 sequence and/or a sequence which specifically hybridizes with the instant  
nucleotide sequence. Hybridizing sequences include similar sequences  
selected from the group of DNA or RNA which specifically interact to the instant  
nucleotide sequences under at least moderate stringency conditions which are  
known in the art. A preferred, non-limiting example of stringent hybridization  
10 conditions are hybridization in 6X sodium chloride/sodium citrate (SSC) at  
about 45°C, followed by one or more washes in 0.2 X SSC, 0.1% SDS at 50-  
65°C. This further includes short nucleotide sequences of e.g. 10 to 30  
nucleotides, preferably 12 to 15 nucleotides. Included are also primer or  
hybridization probes.

15

A homologous nucleotide sequence included within the scope of the instant  
invention is a sequence which is at least about 40%, preferably at least about  
50 % or 60%, and more preferably at least about 70%, 80% or 90% and most  
preferably at least about 95%, 96%, 97%, 98% or 99% or more homologous to  
20 a nucleotide sequence of SEQ ID NO. 1.

All of the aforementioned definitions are true for amino acid sequences and  
functional enzymes and can easily transferred by a person skilled in the art.

Isoenzymes are understood to be enzymes which have the same or a similar  
25 substrate specificity and/or catalytic activity but a different primary structure.

In a first embodiment, this invention is directed to nucleic acid sequences that  
encode a PDAT. This includes sequences that encode biologically active  
PDATs as well as sequences that are to be used as probes, vectors for  
30 transformation or cloning intermediates. The PDAT encoding sequence may

encode a complete or partial sequence depending upon the intended use. All or a portion of the genomic sequence, cDNA sequence, precursor PDAT or mature PDAT is intended.

5 Further included is a nucleotide sequence selected from the group consisting of sequences set forth in SEQ ID No. 1, 1b, 3, 3b, 4, 4a, 4b, 9b, 10, 10b or 11 or a portion, derivate, allele or homolog thereof. The invention pertains a partial nucleotide sequence corresponding to a fulllength nucleotide sequence selected from the group consisting of sequences set forth in SEQ ID No. 5, 5b, 6b, 7, 8b,  
10 9, 11b or 12 or a portion, derivate, allele or homolog thereof. Moreover, a nucleotide sequence comprising a nucleotide sequence which is at least 40% homologous to a nucleotide sequence selected form the group consisting of those sequences set forth in SEQ ID No. 1 1b, 3, 3b, 4, 4a, 4b, 5, 5b, 6b, 7, 8b, 9, 9b, 10, 10b, 11, 11b or 12 is contemplated within the scope of the invention.

15

The instant invention pertains to a gene construct comprising a said nucleotide sequences of the instant invention which is operably linked to a heterologous nucleic acid.

The term operably linked means a serial organisation e.g. of a promotor, coding  
20 sequence, terminator and/or further regulatory elements whereby each element can fulfill its original function during expression of the nucleotide sequence.

Further, a vector comprising of a said nucleotide sequence of the instant invention is contemplated in the instant invention. This includes also an  
25 expression vector as well as a vector further comprising a selectable marker gene and/or nucleotide sequences for the replication in a host cell and/or the integration into the genome of the host cell.

In a different aspect, this invention relates to a method for producing a PDAT in  
30 a host cell or progeny thereof, including genetically engineered oil seeds, yeast and moulds or any other oil accumulating organism, via the expression of a

construct in the cell. Cells containing a PDAT as a result of the production of the PDAT encoding sequence are also contemplated within the scope of the invention.

5 Further, the invention pertains a transgenic cell or organism containing a said nucleotide sequence and/or a said gene construct and/or a said vector. The object of the instant invention is further a transgenic cell or organism which is an eucaryotic cell or organism. Preferably, the transgenic cell or organism is a yeast cell or a plant cell or a plant. The instant invention further pertains said  
10 transgenic cell or organism having an altered biosynthetic pathway for the production of triacylglycerol. A transgenic cell or organism having an altered oil content is also contemplated within the scope of this invention.

Further, the invention pertains a transgenic cell or organism wherein the activity  
15 of PDAT is altered in said cell or organism. This altered activity of PDAT is characterized by an alteration in gene expression, catalytic activity and/or regulation of activity of the enzyme. Moreover, a transgenic cell or organism is included in the instant invention, wherein the altered biosynthetic pathway for the production of triacylglycerol is characterized by the prevention of  
20 accumulation of undesirable fatty acids in the membrane lipids.

In a different embodiment, this invention also relates to methods of using a DNA sequence encoding a PDAT for increasing the oil-content within a cell.

25 Another aspect of the invention relates to the accommodation of high amounts of uncommon fatty acids in the triacylglycerol produced within a cell, by introducing a DNA sequence producing a PDAT that specifically removes these fatty acids from the membrane lipids of the cell and channel them into triacylglycerol. Plant cells having such a modification are also contemplated  
30 herein.

Further, the invention pertains a process for the production of triacylglycerol, comprising growing a said transgenic cell or organism under conditions whereby the said nucleotide sequence is expressed and whereby the said transgenic cells comprising a said enzyme catalysing the transfer of fatty acids from phospholipids to diacylglycerol forming triacylglycerol.

Moreover, triacylglycerols produced by the aforementioned process are included in scope of the instant invention.

Object of the instant invention is further the use of an instant nucleotide sequence and/or a said enzyme for the production of triacylglycerol and/or triacylglycerols with uncommon fatty acids. The use of a said instant nucleotide sequence and/or a said enzyme of the instant invention for the transformation of any cell or organism in order to be expressed in this cell or organism and result in an altered, preferably increased oil content of this cell or organism is also contemplated within the scope of the instant invention.

A PDAT of this invention includes any sequence of amino acids, such as a protein, polypeptide or peptide fragment obtainable from a microorganism, animal or plant source that demonstrates the ability to catalyse the production of triacylglycerol from a phospholipid and diacylglycerol under enzyme reactive conditions. By „enzyme reactive conditions“ is meant that any necessary conditions are available in an environment (e.g., such factors as temperature, pH, lack of inhibiting substances) which will permit the enzyme to function.

Other PDATs are obtainable from the specific sequences provided herein. Furthermore, it will be apparent that one can obtain natural and synthetic PDATs, including modified amino acid sequences and starting materials for synthetic-protein modelling from the exemplified PDATs and from PDATs which are obtained through the use of such exemplified sequences. Modified amino acid sequences include sequences that have been mutated, truncated,

increased and the like, whether such sequences were partially or wholly synthesised. Sequences that are actually purified from plant preparations or are identical or encode identical proteins thereto, regardless of the method used to obtain the protein or sequence, are equally considered naturally  
5 derived.

Further, the nucleic acid probes (DNA and RNA) of the present invention can be used to screen and recover „homologous“ or „related“ PDATs from a variety of plant and microbial sources.

10

Further, it is also apparent that a person skilled in the art can, with the information provided in this application, in any organism identify a PDAT activity, purify an enzyme with this activity and thereby identify a „non-homologous“ nucleic acid sequence encoding such an enzyme.

15

The present invention can be essentially characterized by the following aspects:

1. Use of a PDAT gene (genomic clone or cDNA) for transformation.
- 20 2. Use of a DNA molecule according to item 1 wherein said DNA is used for transformation of any organism in order to be expressed in this organism and result in an active recombinant PDAT enzyme in order to increase oil content of the organism.
3. Use of a DNA molecule of item 1 wherein said DNA is used for  
25 transformation of any organism in order to prevent the accumulation of undesirable fatty acids in the membrane lipids.
4. Use according to item 1, wherein said PDAT gene is used for transforming transgenic oil accumulating organisms engineered to produce any  
uncommon fatty acid which is harmful if present in high amounts in  
30 membrane lipids, such as medium chain fatty acids, hydroxylated fatty acids, epoxygenated fatty acids and acetylenic fatty acids.

5. Use according to item 1, wherein said PDAT gene is used for transforming organisms, and wherein said organisms are crossed with other oil accumulating organisms engineered to produce any uncommon fatty acid which is harmful if present in high amounts in membrane lipids, comprising medium chain fatty acids, hydroxylated fatty acids, epoxygenated fatty acids and acetylenic fatty acids.
6. Use according to item 1, wherein the enzyme encoded by said PDAT gene or cDNA is coding for a PDAT with distinct acyl specificity.
7. Use according to item 1 wherein said PDAT encoding gene or cDNA, is derived from *Saccharomyces cerevisiae*, or contain nucleotide sequences coding for an amino acid sequence 30% or more identical to the amino acid sequence of PDAT as presented in SEQ. ID. NO. 2.
8. Use according to item 1 wherein said PDAT encoding gene or cDNA is derived from *Saccharomyces cerevisiae*, or contain nucleotide sequences coding for an amino acid sequence 40% or more *identical* to the amino acid sequence of PDAT as presented in SEQ. ID. NO. 2.
9. Use according to item 1 wherein said PDAT encoding gene or cDNA is derived from *Saccharomyces cerevisiae*, or contain nucleotide sequences coding for an amino acid sequence 60% or more *identical* to the amino acid sequence of PDAT as presented in SEQ. ID. NO. 2.
10. Use according to item 1 wherein said PDAT encoding gene or cDNA is derived from *Saccharomyces cerevisiae*, or contain nucleotide sequences coding for an amino acid sequence 80% or more identical to the amino acid sequence of PDAT as presented in SEQ. ID. NO. 2.
11. Use according to item 1 wherein said PDAT encoding gene or cDNA is derived from plants or contain nucleotide sequences coding for an amino acid sequence 40% or more identical to the amino acid sequence of PDAT from *Arabidopsis thaliana* or to the protein encoded by the fulllength counterpart of the partial Zea mays, Lycopericon esculentum, or Neurospora crassa cDNA clones.

12. Transgenic oil accumulating organisms comprising, in their genome, a PDAT gene transferred by recombinant DNA technology or somatic hybridization.
13. Transgenic oil accumulating organisms according to item 12 comprising, in  
5 their genome, a PDAT gene having specificity for substrates with a particular uncommon fatty acid and the gene for said uncommon fatty acid.
14. Transgenic organisms according to item 12 or 13 which are selected from the group consisting of fungi, plants and animals.
15. Transgenic organisms according to item 12 or 13 which are selected from  
10 the group of agricultural plants.
16. Transgenic organisms according to item 12 or 13 which are selected from the group of agricultural plants and where said PDAT gene is expressed under the control of a storage organ specific promotor.
17. Transgenic organisms according to item 12 or 13 which are selected from  
15 the group of agricultural plants and where said PDAT gene is expressed under the control of a seed promotor.
18. Oils from organisms according to item 12 – 17.
19. A method for altering acyl specificity of a PDAT by alteration of the nucleotide sequence of a naturally occurring encoding gene and as a  
20 consequence of this alternation creating a gene encoding for an enzyme with novel acyl specificity.
20. A protein encoded by a DNA molecule according to item 1 or a functional fragment thereof.
21. A protein of item 20 designated phospholipid:diacylglycerol acyltransferase.
- 25 22. A protein of item 21 which has a distinct acyl specificity.
23. A protein of item 13 having the amino acid sequence as set forth in SEQ, ID NO. 2, 13, 14 or 15 (and the proteins encoded by the fulllength or partial genes set forth in SEQ. ID. NO. 1, 3, 4, 5, 7, 9, 10, 11 or 12) or an amino acid sequence with at least 30 % homology to said amino acid sequence.
- 30 24. A protein of item 23 isolated from *Saccharomyces cerevisiae*.



General methods:

Yeast strains and plasmids. The wild type yeast strains used were either FY1679 (*MAT $\alpha$  his3- $\Delta$ 200 leu2- $\Delta$ 1 trp1- $\Delta$ 6 ura3-52*) or W303-1A (*MAT $\alpha$  ADE2-1 can1-100 his3-11,15 leu2-3,112 trp1-1 ura3-1*) (7). The YNR008w::KanMX2 disruption strain FVKT004-04C(AL), which is congenic to FY1679, was obtained from the Euroscarf collection (8). A 2751 bp fragment containing the YNR008w gene with 583 bp of 5' and 183 bp of 3' flanking DNA was amplified from W303-1A genomic DNA using *Taq* polymerase with 5'-TCTCCATCTTCTGCAAAACCT-3' and 5'-CCTGTCAAAAACCTTCTCCTC-3' as primers. The resulting PCR product was purified by agarose gel electrophoresis and cloned into the *EcoRV* site of pBluescript (pbluescript-pdat). For complementation experiments, the cloned fragment was released from pBluescript by *HindIII*-*SacI* digestion and then cloned between the *HindIII* and *SacI* sites of pFL39 (9), thus generating pUS1. For overexpression of the PDAT gene, a 2202 bp *EcoRI* fragment from the pBluscript plasmid which contains only 24 bp of 5' flanking DNA was cloned into the *BamHI* site of the *GAL1-TPK2* expression vector pJN92 (12), thus generating pUS4.

Microsomal preparations. Microsomes from developing seeds of sunflower (*Helianthus annuus*), *Ricinus communis* and *Crepis palaestina* were prepared using the procedure of Stobart and Stymne (11). To obtain yeast microsomes, 1g of yeast cells (fresh weight) was re-suspended in 8 ml of ice-cold buffer (20 mM Tris-Cl, pH 7.9, 10 mM MgCl<sub>2</sub>, 1 mM EDTA, 5 % (v/v) glycerol, 1 mM DTT, 0.3 M ammonium sulfate) in a 12 ml glass tube. To this tube, 4 ml of glass beads (diameter 0.45-0.5 mm) were added, and the tube was then heavily shaken (3 x 60 s) in an MSK cell homogenizer (B. Braun Melsungen AG, Germany). The homogenized suspension was centrifuged at 20,000 x g for 15 min at 6°C and the resulting supernatant was again centrifuged at 100,000 x g for 2 h at 6°C. The 100,000 x g pellet was resuspended in 0.1 M potassium

phosphate (pH 7.2), and stored at -80°C. It is subsequently referred to as the crude yeast microsomal fraction.

Lipid substrates. Radio-labeled ricinoleic (12-hydroxy-9-octadecenoic) and vernolic (12,13-epoxy-9-octadecenoic) acids were synthesized enzymatically from [ $^{14}\text{C}$ ]oleic acid and [ $^{14}\text{C}$ ]linoleic acid, respectively, by incubation with microsomal preparations from seeds of *Ricinus communis* and *Crepis palaestina*, respectively (12). The synthesis of phosphatidylcholines (PC) or phosphatidylethanolamines (PE) with  $^{14}\text{C}$ -labeled acyl groups in the *sn*-2 position was performed using either enzymatic (13), or synthetic (14) acylation of [ $^{14}\text{C}$ ]oleic, [ $^{14}\text{C}$ ]ricinoleic, or [ $^{14}\text{C}$ ]vernolic acid. Dioleoyl-PC that was labeled in the *sn*-1 position was synthesized from *sn*-1-[ $^{14}\text{C}$ ]oleoyl-lyso-PC and unlabeled oleic acid as described in (14). *Sn*-1-oleoyl-*sn*-2-[ $^{14}\text{C}$ ]ricinoleoyl-DAG was synthesized from PC by the action of phospholipase C type XI from *B. Cereus* (Sigma Chemical Co.) as described in (15). Monovernoloyl- and divernoloyl-DAG were synthesized from TAG extracted from seeds of *Euphorbia lagascae*, using the TAG-lipase (*Rizhopus arrhizus*, Sigma Chemical Co.) as previously described (16). Monoricinoleoyl-TAG was synthesized according to the same method using TAG extracted from Castor bean.

20

Lipid analysis. Total lipid composition of yeast were determined from cells harvested from a 40 ml liquid culture, broken in a glass-bead shaker and extracted into chloroform as described by Bligh and Dyer (17), and then separated by thin layer chromatography in hexane/diethylether/acetic acid (80:20:1) using pre-coated silica gel 60 plates (Merck). The lipid areas were located by brief exposure to  $\text{I}_2$  vapors and identified by means of appropriate standards. Polar lipids, sterol-esters and triacylglycerols, as well as the remaining minor lipid classes, referred to as other lipids, were excised from the plates. Fatty acid methylesters were prepared by heating the dry excised material at 85 °C for 60 min in 2% (v/v) sulfuric acid in dry methanol. The methyl esters were extracted with hexane and analyzed by GLC through a 50 m

30

x 0.32 mm CP-Wax58-CB fused-silica column (Chrompack), with methylheptadecanoic acid as an internal standard. The fatty acid content of each fraction was quantified and used to calculate the relative amount of each lipid class. In order to determine the total lipid content, 3 ml aliquots from yeast  
5 cultures were harvested by centrifugation and the resulting pellets were washed with distilled water and lyophilized. The weight of the dried cells was determined and the fatty acid content was quantified by GLC-analyses after conversion to methylesters as described above. The lipid content was then calculated as nmol fatty acid (FA) per mg dry weight yeast.

10

Enzyme assays. Aliquots of crude microsomal fractions (corresponding to 10 nmol of microsomal PC) from developing plant seeds or yeast cells were lyophilized over night. <sup>14</sup>C-Labeled substrate lipids dissolved in benzene were then added to the dried microsomes. The benzene was evaporated under a  
15 stream of N<sub>2</sub>, leaving the lipids in direct contact with the membranes, and 0.1 ml of 50 mM potassium phosphate (pH 7.2) was added. The suspension was thoroughly mixed and incubated at 30°C for the time period indicated, up to 90 min. Lipids were extracted from the reaction mixture using chloroform and separated by thin layer chromatography in hexane/diethylether/acetic acid  
20 (35:70:1.5) using silica gel 60 plates (Merck). The radioactive lipids were visualized and quantified on the plates by electronic autoradiography (Instant Imager, Packard, US).

Yeast cultivation. Yeast cells were grown at 28°C on a rotatory shaker in  
25 liquid YPD medium (1% yeast extract, 2% peptone, 2% glucose), synthetic medium (18) containing 2% (v/v) glycerol and 2% (v/v) ethanol, or minimal medium (19) containing 16 g/l of glycerol.

The instant invention is further characterized by the following examples which  
30 are not limiting:

Acyl-CoA-independent synthesis of TAG by oil seed microsomes. A large number of unusual fatty acids can be found in oil seeds (20). Many of these fatty acids, such as ricinoleic (21) and vernolic acids (22), are synthesized using phosphatidylcholin (PC) with oleoyl or linoleoyl groups esterified to the *sn*-2 position, respectively, as the immediate precursor. However, even though PC can be a substrate for unusual fatty acid synthesis and is the major membrane lipids in seeds, unusual fatty acids are rarely found in the membranes. Instead, they are mainly incorporated into the TAG. A mechanism for efficient and selective transfer of these unusual acyl groups from PC into TAG must therefore exist in oil seeds that accumulate such unusual fatty acids. This transfer reaction was biochemically characterized in seeds from castor bean (*Ricinus communis*) and *Crepis palaestina*, plants which accumulate high levels of ricinoleic and vernolic acid, respectively, and sunflower (*Helianthus annuus*), a plant which has only common fatty acids in its seed oil. Crude microsomal fractions from developing seeds were incubated with PC having <sup>14</sup>C-labeled oleoyl, ricinoleoyl or vernoloyl groups at the *sn*-2 position. After the incubation, lipids were extracted and analyzed by thin layer chromatography. We found that the amount of radioactivity that was incorporated into the neutral lipid fraction increased linearly over a period of 4 hours (data not shown). The distribution of [<sup>14</sup>C]acyl groups within the neutral lipid fraction was analyzed after 80 min (Fig. 1). Interestingly the amount and distribution of radioactivity between different neutral lipids were strongly dependent both on the plant species and on the type of [<sup>14</sup>C]acyl chain. Thus, sunflower microsomes incorporated most of the label into DAG, regardless of the type of [<sup>14</sup>C]acyl group. In contrast, *R. communis* microsomes preferentially incorporated [<sup>14</sup>C]ricinoleoyl and [<sup>14</sup>C]vernoloyl groups into TAG, while [<sup>14</sup>C]oleyl groups mostly were found in DAG. *C. palaestina* microsomes, finally, incorporated only [<sup>14</sup>C]vernoloyl groups into TAG, with [<sup>14</sup>C]ricinoleyl groups being found mostly as free fatty acids, and [<sup>14</sup>C]oleyl groups in DAG. This shows that the high *in vivo* levels of ricinoleic acid and vernolic acid in the TAG pool of *R. communis*

and *C. palaestina*, respectively, can be explained by an efficient and selective transfer of the corresponding acyl groups from PC to TAG in these organisms.

5 The in-vitro synthesis of triacylglycerols in microsomal preparations of developing castor bean is summarized in table 1.

PDAT: a novel enzyme that catalyzes acyl-CoA independent synthesis of TAG. It was investigated if DAG could serve both as an acyl donor as well as an acyl acceptor in the reactions catalyzed by the oil seed microsomes. 10 Therefore, unlabeled divernoloyl-DAG was incubated with either *sn*-1-oleoyl-*sn*-2-[<sup>14</sup>C]ricinoleoyl-DAG or *sn*-1-oleoyl-*sn*-2-[<sup>14</sup>C]ricinoleoyl-PC in the presence of *R. communis* microsomes. The synthesis of TAG molecules containing both [<sup>14</sup>C]ricinoleoyl and vernoloyl groups was 5 fold higher when [<sup>14</sup>C]ricinoleoyl-PC served as acyl donor as compared to [<sup>14</sup>C]ricinoleoyl-DAG 15 (fig.1B). These data strongly suggests that PC is the immediate acyl donor and DAG the acyl acceptor in the acyl-CoA-independent formation of TAG by oil seed microsomes. Therefore, this reaction is catalyzed by a new enzyme which we call phospholipid : diacylglycerol acyltransferase (PDAT).

20 PDAT activity in yeast microsomes. Wild type yeast cells were cultivated under conditions where TAG synthesis is induced. Microsomal membranes were prepared from these cells and incubated with *sn*-2-[<sup>14</sup>C]-ricinoleoyl-PC and DAG and the <sup>14</sup>C-labeled products formed were analyzed. The PC-derived [<sup>14</sup>C]ricinoleoyl groups within the neutral lipid fraction mainly were found in free 25 fatty acids or TAG, and also that the amount of TAG synthesized was dependent on the amount of DAG that was added to the reaction (Fig.2). The *in vitro* synthesis of TAG containing both ricinoleoyl and vernoloyl groups, a TAG species not present *in vivo*, from exogenous added *sn*-2-[<sup>14</sup>C]ricinoleoyl-PC and unlabelled vernoloyl-DAG (Fig. 2, lane 3) clearly demonstrates the existence of 30 an acyl-CoA-independent synthesis of TAG involving PC and DAG as

substrates in yeast microsomal membranes. Consequently, TAG synthesis in yeast can be catalyzed by an enzyme similar to the PDAT found in plants.

The PDAT encoding gene in yeast.

5 A gene in the yeast genome (YNR008w) is known, but nothing is known about the function of YNR008w, except that the gene is not essential for growth under normal circumstances. Microsomal membranes were prepared from the yeast strain FVKT004-04C(AL) (8) in which this gene with unknown function had been disrupted. PDAT activity in the microsomes were assayed using PC with  
10 radiolabelled fatty acids at the sn-2 position. The activity was found to be completely absent in the disruption strain (Fig. 2 lane 4). Significantly, the activity could be partially restored by the presence of YNR008w on the single copy plasmid pUS1 (Fig. 2 lane 5). Moreover, acyl groups of phosphatidylethanolamine (PE) were efficiently incorporated into TAG by  
15 microsomes from the wild type strain whereas no incorporation occurred from this substrate in the mutant strain (data not shown). This shows that YNR008w encodes a yeast PDAT which catalyzes the transfer of an acyl group from the sn-2 position of phospholipids to DAG, thus forming TAG. It should be noted that no cholesterol esters were formed from radioactive PC even in incubations  
20 with added ergosterols, nor were the amount of radioactive free fatty acids formed from PC affected by disruption of the YNR008w gene (data not shown). This demonstrates that yeast PDAT do not have cholesterol ester synthesising or phospholipase activities.

25 Increased TAG content in yeast cells that overexpress PDAT. The effect of overexpressing the PDAT-encoding gene was studied by transforming a wild type yeast strain with the pUS4 plasmid in which the gene is expressed from the galactose-induced *GAL1:TPK2* promoter. Cells containing the empty expression vector were used as a control. The cells were grown in synthetic  
30 glycerol-ethanol medium, and expression of the gene was induced after either 2 hours (early log phase) or 25 hours (stationary phase) by the addition of

galactose. The cells were then incubated for another 21 hours, after which they were harvested and assays were performed. We found that overexpression of PDAT had no significant effect on the growth rate as determined by the optical density. However, the total lipid content, measured as  $\mu\text{mol}$  fatty acids per mg yeast dry weight, was 47% (log phase) or 29% (stationary phase) higher in the PDAT overexpressing strain than in the control. Furthermore, the polar lipid and sterolester content was unaffected by overexpression of PDAT. Instead, the elevated lipid content in these cells is entirely due to an increased TAG content (Fig. 3A,B). Thus, the amount of TAG was increased by 2-fold in PDAT overexpressing early log phase cells and by 40% in stationary phase cells. It is interesting to note that a significant increase in the TAG content was achieved by overexpressing PDAT even under conditions (*i.e.* in stationary phase) where DAGAT is induced and thus contributes significantly to TAG synthesis. *In vitro* PDAT activity assayed in microsomes from the PDAT overexpressing strain was 7-fold higher than in the control strain, a finding which is consistent with the increased levels of TAG that we observed *in vivo* (Fig. 3C). These results clearly demonstrate the potential use of the PDAT gene in increasing the oil content in transgenic organisms.

Substrate specificity of yeast PDAT. The substrate specificity of yeast PDAT was analyzed using microsomes prepared from the PDAT overexpressing strain (see Fig. 4). The rate of TAG synthesis, under conditions given in figure 4 with di-oleoyl-PC as the acyl-donor, was 0.15 nmol per min and mg protein. With both oleoyl groups of PC labeled it was possible, under the given assay conditions, to detect the transfer of 11 pmol/min of [ $^{14}\text{C}$ ]oleoyl chain into TAG and the formation of 15 pmol/min of lyso-PC. In microsomes from the PDAT-deficient strain, no TAG at all and only trace amounts of lyso-PC was detected, strongly suggesting that yeast PDAT catalyses the formation of equimolar amounts of TAG and lyso-PC when supplied with PC and DAG as substrates. The fact that somewhat more lyso-PC than TAG is formed can be

explained by the presence of a phospholipase in yeast microsomes, which produces lyso-PC and unesterified fatty acids from PC.

The specificity of yeast PDAT for different acyl group positions was investigated by incubating the microsomes with di-oleoyl-PC carrying a [<sup>14</sup>C]acyl group either at the *sn*-1 position (Fig. 4A bar 2) or the *sn*-2 position (Fig. 4A bar 3). We found that the major <sup>14</sup>C-labeled product formed in the former case was lyso-PC, and in the latter case TAG. We conclude that yeast PDAT has a specificity for the transfer of acyl groups from the *sn*-2 position of the phospholipid to DAG, thus forming *sn*-1-lyso-PC and TAG. Under the given assay conditions, trace amounts of <sup>14</sup>C-labelled DAG is formed from the *sn*-1 labeled PC by the reversible action of a CDP-choline : choline phosphotransferase. This labeled DAG can then be further converted into TAG by the PDAT activity. It is therefore not possible to distinguish whether the minor amounts of labeled TAG that is formed in the presence of di-oleoyl-PC carrying a [<sup>14</sup>C]acyl group in the *sn*-1 position, is synthesized directly from the *sn*-1-labeled PC by a PDAT that also can act on the *sn*-1 position, or if it is first converted to *sn*-1-labeled DAG and then acylated by a PDAT with strict selectivity for the transfer of acyl groups at the *sn*-2 position of PC. Taken together, this shows that the PDAT encoded by YNR008w catalyses an acyl transfer from the *sn*-2 position of PC to DAG, thus causing the formation of TAG and lyso-PC.

The substrate specificity of yeast PDAT was further analyzed with respect to the headgroup of the acyl donor, the acyl group transferred and the acyl chains of the acceptor DAG molecule. The two major membrane lipids of *S. cerevisiae* are PC and PE, and as shown in Fig. 4B (bars 1 and 2), dioleoyl-PE is nearly 4-fold more efficient than dioleoyl-PC as acyl donor in the PDAT-catalyzed reaction. Moreover, the rate of acyl transfer is strongly dependent on the type of acyl group that is transferred. Thus, a ricinoleoyl group at the *sn*-2 position of PC is 2.5 times more efficiently transferred into TAG than an oleoyl



group in the same position (Fig. 4B bars 1 and 3). In contrast, yeast PDAT has no preference for the transfer of vernoloyl groups over oleoyl groups (Fig. 4B bars 1 and 4). The acyl chain of the acceptor DAG molecule also affects the efficiency of the reaction. Thus, DAG with a ricinoleoyl or a vernoloyl group is a more efficient acyl acceptor than dioleoyl-DAG (Fig. 4B bars 1, 5 and 6). Taken together, these results clearly show that the efficiency of the PDAT-catalyzed acyl transfer is strongly dependent on the properties of the substrate lipids.

PDAT genes. Nucleotide and amino acid sequences of several PDAT genes are given as SEQ ID No. 1 through 15. Further provisional and/or partial sequences are given as SEQ ID NO 1a through 5a and 1b through 11b, respectively. One of the Arabidopsis genomic sequences (SEQ ID NO. 4) identified an Arabidopsis EST cDNA clone; T04806. This cDNA clone was fully characterised and the nucleotide sequence is given as SEQ ID NO. 5. Based on the sequence homology of the T04806 cDNA and the *Arabidopsis thaliana* genomic DNA sequence (SEQ ID NO 4) it is apparent that an additional A is present at position 417 in the cDNA clone (data not shown). Excluding this nucleotide would give the amino acid sequence depicted in SEQ ID NO. 12.

Increased TAG content in seeds of Arabidopsis thaliana that express the yeast PDAT. For the expression of the yeast PDAT gene in *Arabidopsis thaliana* an EcoRI fragment from the pBluescript-PDAT was cloned together with napin promotor (25) into the vector pGPTV-KAN (26). A plasmid (pGNapPDAT) having the yeast PDAT gene in the correct orientation was identified and transformed into *Agrobacterium tumefaciens*. These bacteria were used to transform *Arabidopsis thaliana* columbia (C-24) plants using the root transformation method (27). Plants transformed with an empty vector were used as controls.

First generation seeds (T1) were harvested and germinated on kanamycin containing medium. Second generation seeds (T2) were pooled from individual plants and their fatty acid contents analysed by quantification of their methyl

esters by gas liquid chromatography after methylation of the seeds with 2% sulphuric acid in methanol at 85 °C for 1,5 hours. Quantification was done with heptadecanoic acid methyl esters as internal standard.

5 From the transformation with pGNapPDAT one T1 plant (26-14) gave raise to seven T2 plants of which 3 plants yielded seeds with statistically (in a mean difference two-sided test) higher oil content than seeds from T2 plants generated from T1 plant 32-4 transformed with an empty vector (table 2).

## References cited in the description:

1. Bell, R. M. & Coleman, R. A. (1980) *Annu. Rev. Biochem.* **49**, 459-487.
2. Stymne, S. & Stobart, K. (1987) in *The biochemistry of plants: a comprehensive treatise*, Vol. 9, ed. Stumpf, P. K. (Academic Press, New York), pp. 175-214.
3. Cases, S. et al. (1998) *Proc. Natl. Acad. Sci. U S A* **95**, 13018-13023.
4. Hobbs, D. H., Lu, C. & Hills, M. J. (1999) *FEBS Lett.* **452**, 145-9
5. Zou, J., Wei, Y., Jako, C., Kumar, A., Selvaraj, G. & Taylor, D. C. (1999) *Plant J.* **19**, 645-653.
6. Lardizabal, K., Hawkins, D., Mai, J., & Wagner, N. (1999) Abstract presented at the Biochem. Mol. Plant Fatty Acids Glycerolipids Symposium, South Lake Tahoe, USA.
7. Thomas, B. J. & Rothstein, R. (1989) *Cell* **56**, 619-630.
8. Entian, K.-D. & Kötter, P. (1998) *Meth. Microbiol.* **26**, 431-449.
9. Kern, L., de Montigny, J., Jund, R. & Lacroute, F. (1990) *Gene* **88**, 149-157.
10. Ronne, H., Carlberg, M., Hu, G.-Z. & Nehlin, J. O. (1991) *Mol. Cell. Biol.* **11**, 4876-4884.
11. Stobart, K. & Stymne, S. (1990) in *Method in Plant Biochemistry*, vol 4, eds. Harwood, J. L. & Bowyer, J. R. (Academic press, London), pp. 19-46.
12. Bafor, M., Smith, M. A., Jonsson, L., Stobrt, A. K. & Stymne, S. (1991) *Biochem. J.* **280**, 507-514.
13. Banas, A., Johansson, I. & Stymne, S. (1992) *Plant Science* **84**, 137-144.
14. Kanda, P. & Wells, M. A. (1981) *J. Lipid. Res.* **22**, 877-879.
15. Ståhl, U., Ek, B. & Stymne, S. (1998) *Plant Physiol.* **117**, 197-205.
16. Stobart, K., Mancha, M. & Lenman M, Dahlqvist, A. & Stymne, S. (1997) *Planta* **203**, 58-66.
17. Bligh, E. G. & Dyer, W. J. (1959) *Can. J. Biochem. Physiol.* **37**, 911-917.
18. Sherman, F., Fink, G. R. & Hicks, J. B. (1986) in *Laboratory Course Manual for Methods in Yeast Genetics* (Cold Spring Harbor Laboratory)
19. Meesters, P. A. E. P., Huijberts, G. N. M. and Eggink, G. (1996) *Appl. Microbiol. Biotechnol.* **45**, 575-579.
20. van de Loo, F. J., Fox, B. G. & Sommerville, C. (1993), in *Lipid metabolism in plants*, ed. Moore, T. S. (CRC Press, Inc.), pp. 91-126.
21. van de Loo, F. J., Broun, P., Turner, S. & Sommerville, S. (1995) *Proc. Natl.*

*Acad. Sci. U S A* **95**, 6743-6747.

22. Lee, M., Lenman, M., Banas, A., Bafor, M., Singh, S., Schweizer, M., Nilsson, R., Liljenberg, C., Dahlqvist, A., Gummesson, P-O., Sjö Dahl, S., Green, A., and Stymne, S. (1998) *Science* **280**, 915-918.
- 5 23. Thompson, J. D., Gibson, T. J., Plewniak, F., Jeanmougin, F. & Higgins, D. G. (1997) *Nucl. Acids Res.* **24**, 4876-4882.
24. Saitou, N. & Nei, M. (1987) *Mol. Biol. Evol.* **4**, 406-425.
25. Stålberg, K., Ellerström, M., Josefsson, L., & Rask, L. (1993) *Plant Mol. Biol.* **23**, 671
- 10 26. Becker, D., Kemper, E., Schell, J., Masterson, R. (1992) *Plant Mol. Biol.* **20**, 1195
27. D. Valvekens, M. Van Montagu, and Van Lusbettens (1988) *Proc. Natl. Acad. Sci. U.S.A.* **85**, 5536

15

## Description of Figures

FIG. 1.

Metabolism of  $^{14}\text{C}$ -labeled PC into the neutral lipid fraction by plant microsomes. (A) Microsomes from developing seeds of sunflower, *R. communis* and *C. palaestina* were incubated for 80 min at 30°C with PC (8 nmol) having oleic acid in its *sn*-1 position, and either  $^{14}\text{C}$ -labeled oleic, ricinoleic or vernolic acid in its *sn*-2 position. Radioactivity incorporated in TAG (open bars), DAG (solid bars), and unsterified fatty acids (hatched bars) was quantified using thin layer chromatography followed by electronic autoradiography, and is shown as percentage of added labeled substrate. (B) Synthesis *in vitro* of TAG carrying two vernoloyl and one [ $^{14}\text{C}$ ]ricinoleoyl group by microsomes from *R. communis*. The substrates added were unlabeled divernoloyl-DAG (5 nmol), together with either *sn*-1-oleoyl-*sn*-2-[ $^{14}\text{C}$ ]ricinoleoyl-DAG (0.4 nmol, 7700 dpm/nmol) or *sn*-1-oleoyl-*sn*-2-[ $^{14}\text{C}$ ]ricinoleoyl-PC (0.4 nmol, 7700 dpm/nmol). The microsomes were incubated with the substrates for 30 min at 30°C, after which samples were removed for lipid analysis as described in the section „general methods“. The data shown are the average of two experiments.

FIG. 2.

PDAT activity in yeast microsomes, as visualized by autoradiogram of neutral lipid products separated on TLC. Microsomal membranes (10 nmol of PC) from the wild type yeast strain FY1679 (lanes 1-3), a congenic yeast strain (FVKT004-04C(AL)) that is disrupted for YNR008w (lane 4) or the same disruption strain transformed with the plasmid pUS1, containing the YNR008w gene behind its native promotor (lane 5), were assayed for PDAT activity. As substrates, we used 2 nmol *sn*-1-oleoyl-*sn*-2-[ $^{14}\text{C}$ ]ricinoleoyl-PC together with either 5 nmol of dioleoyl-DAG (lanes 2, 4 and 5) or *rac*-oleoyl-vernoleoyl-DAG (lane 3). The enzymatic assay and lipid analysis was performed as described in Materials and Methods. The cells were precultured for 20 h in liquid YPD

medium, harvested and re-suspended in an equal volume of minimal medium (19) containing 16 g/l glycerol. The cells were then grown for an additional 24 h prior to being harvested. Selection for the plasmid was maintained by growing the transformed cells in synthetic medium lacking uracil (18). Abbreviations: 1-OH-TAG, monoricinoleoyl-TAG; 1-OH-1-ep-TAG, monoricinoleoyl-monovernoloyl-TAG; OH-FA, unesterified ricinoleic acid.

Fig. 3.

Lipid content (A,B) and PDAT activity (C) in PDAT overexpressing yeast cells.

- 10 The PDAT gene in the plasmid pUS4 was overexpressed from the galactose-induced *GAL1-TPK2* promoter in the wild type strain W303-1A (7). Its expression was induced after (A) 2 hours or (B) 25 hours of growth by the addition of 2% final concentration (w/v) of galactose. The cells were then incubated for another 22 hours before being harvested. The amount of lipids of
- 15 the harvested cells was determined by GLC-analysis of its fatty acid contents and is presented as  $\mu\text{mol}$  fatty acids per mg dry weight in either TAG (open bar), polar lipids (hatched bar), sterol esters (solid bar) and other lipids (striped bar). The data shown are the mean values of results with three independent yeast cultures. (C) *In vitro* synthesis of TAG by microsomes prepared from
- 20 yeast cells containing either the empty vector (vector) or the PDAT plasmid (+ PDAT). The cells were grown as in Fig. 3A. The substrate lipids dioleoyl-DAG (2.5 nmol) and *sn*-1-oleoyl-*sn*-2-[ $^{14}\text{C}$ ]-oleoyl-PC (2 nmol) were added to aliquots of microsomes (10 nmol PC), which were then incubated for 10 min at 28 °C. The amount of label incorporated into TAG was quantified by electronic
- 25 autoradiography. The results shown are the mean values of two experiments.

FIG. 4.

- Substrate specificity of yeast PDAT. The PDAT activity was assayed by incubating aliquots of lyophilized microsomes (10 nmol PC) with substrate lipids
- 30 at 30°C for 10 min (panel A) or 90 min (panel B). Unlabeled DAG (2.5 nmol) was used as substrates together with different labeled phospholipids, as shown

in the figure. (A) *Sn*-position specificity of yeast PDAT regarding the acyl donor substrate. Dioleoyl-DAG together with either *sn*-1-[<sup>14</sup>C]oleoyl-*sn*-2-[<sup>14</sup>C]oleoyl-PC (di-[<sup>14</sup>C]-PC), *sn*-1-[<sup>14</sup>C]oleoyl-*sn*-2-oleoyl-PC (*sn*1-[<sup>14</sup>C]-PC) or *sn*-1-oleoyl-*sn*-2-[<sup>14</sup>C]oleoyl-PC (*sn*2-[<sup>14</sup>C]-PC). (B) Specificity of yeast PDAT regarding  
5 phospholipid headgroup and of the acyl composition of the phospholipid as well as of the diacylglycerol. Dioleoyl-DAG together with either *sn*-1-oleoyl-*sn*-2-[<sup>14</sup>C]oleoyl-PC (oleoyl-PC), *sn*-1-oleoyl-*sn*-2-[<sup>14</sup>C]oleoyl-PE (oleoyl-PE), *sn*-1-oleoyl-*sn*-2-[<sup>14</sup>C]ricinoleoyl-PC (ricinoleoyl-PC) or *sn*-1-oleoyl-*sn*-2-[<sup>14</sup>C]vernoloyl-PC (vernoloyl-PC). In the experiments presented in the 2 bars to  
10 the far right, monoricinoleoyl-DAG (ricinoleoyl-DAG or mono-vernoloyl-DAG (vernoloyl-DAG) were used together with *sn*-1-oleoyl-*sn*-2-[<sup>14</sup>C]-oleoyl-PC. The label that was incorporated into TAG (solid bars) and lyso-PC (LPC, open bars) was quantified by electronic autoradiography. The results shown are the mean values of two experiments. The microsomes used were from W303-1A cells  
15 overexpressing the PDAT gene from the *GAL1-TPK2* promotor, as described in Fig. 3. The expression was induced at early stationary phase and the cells were harvested after an additional 24 h.

20 TAB.1:

In vitro synthesis of triacylglycerols in microsomal preparations of developing castor bean. Aliquots of microsomes (20 nmol PC) were lyophilised and substrate lipids were added in benzene solution: (A) 0.4 nmol [<sup>14</sup>C]-DAG (7760 dpm/nmol) and where indicated 1.6 nmol unlabelled DAG; (B) 0.4 nmol [<sup>14</sup>C]-  
25 DAG (7760 dpm/nmol) and 5 nmol unlabelled di-ricinoleoyl-PC and (C) 0.25 nmol [<sup>14</sup>C]-PC (4000 dpm/nmol) and 5 nmol unlabelled DAG. The benzene was evaporated by N<sub>2</sub> and 0.1 ml of 50 mM potassium phosphate was added, thoroughly mixed and incubated at 30 °C for (A) 20 min.; (B) and (C) 30 min.. Assays were terminated by extraction of the lipids in chloroform. The lipids  
30 were then separated by thin layer chromatography on silica gel 60 plates

(Merck; Darmstadt, Germany) in hexan/diethylether/acetic 35:70:1.5. The radioactive lipids were visualised and the radioactivity quantified on the plate by electronic autoradiography (Instant Imager, Packard, US). Results are presented as mean values of two experiments.

5

Radioactivity in different triacylglycerols (TAG) species formed. Abbreviations used: 1-OH-, mono-ricinoleoyl-; 2-OH, di-ricinoleoyl-; 3-OH-, triricinoleoyl-; 1-OH-1-ver-, mono-ricinoleoyl-monovernoleoyl-; 1-OH-2-ver-, mono-ricinoleoyl-divernoleoyl-. Radiolabelled DAG and PC were prepared enzymatically. The radiolabelled ricinoleoyl group is attached at the sn-2-position of the lipid and unlabelled oleoyl group at the sn-1-position. Unlabelled DAG with vernoleoyl- or ricinoleoyl chains were prepared by the action of TAG lipase (6) on oil of *Euphorbia lagascae* or Castor bean, respectively. Synthetic di-ricinoleoyl-PC was kindly provided from Metapontum Agribios (Italy).

15

#### TAB.2:

Total fatty acids per mg of T2 seeds pooled from individual *Arabidopsis thaliana* plants transformed with yeast PDAT gene under the control of napin promotor (26-14) or transformed with empty vector (32-4).

20

\* = statistical difference between control plants and PDAT transformed plants in a mean difference two-sided test at  $\alpha = 5$ .



Description of the SEQ ID:

5     *SEQ ID NO. 1:* Genomic DNA sequence and suggested amino acid sequence of the *Saccharomyces cerevisiae* PDAT gene, YNR008w, with GenBank accession number Z71623 and Y13139, and with nucleotide ID number 1302481.

*SEQ ID NO. 2:* The amino acid sequence of the suggested open reading frame YNR008w from *Saccharomyces cerevisiae*.

10    *SEQ ID NO. 3:* Genomic DNA sequence of the *Schizosaccharomyces pombe* gene SPBC776.14.

*SEQ ID NO. 4:* Genomic DNA sequence of part of the *Arabidopsis thaliana* locus with GenBank accession number AB006704.

15    *SEQ ID NO. 5:* Nucleotide sequence of the *Arabidopsis thaliana* cDNA clone with GenBank accession number T04806, and nucleotide ID number 315966.

*SEQ ID NO. 6:* Predicted amino acid sequence of the *Arabidopsis thaliana* cDNA  
20    clone with GenBank accession number T04806.

*SEQ ID NO. 7:* Nucleotide and amino acid sequence of the *Zea mays* EST clone with GenBank accession number AI491339, and nucleotide ID number 4388167.

25    *SEQ ID NO. 8:* Predicted amino acid sequence of the *Zea mays* EST clone with GenBank accession number AI491339, and nucleotide ID number 4388167.

*SEQ ID NO. 9:* DNA sequence of part of the *Neurospora crassa* EST clone W07G1, with GenBank accession number AI398644, and nucleotide ID number  
30    4241729.

SEQ ID NO. 10: Genomic DNA sequence of part of the *Arabidopsis thaliana* locus with GenBank accession number AC004557.

5 SEQ ID NO. 11: Genomic DNA sequence of part of the *Arabidopsis thaliana* locus with GenBank accession number AC003027.

SEQ ID NO. 12: DNA sequence of part of the *Lycopersicon esculentum* cDNA clone with GenBank accession number AI486635.

10 SEQ ID NO. 13: Amino acid sequence of the *Schizosaccharomyces pombe* putative open reading frame CAA22887 of the *Schizosaccharomyces pombe* gene SPBC776.14.

15 SEQ ID NO. 14: Amino acid sequence of the *Arabidopsis thaliana* putative open reading frame AAC80628 derived from the *Arabidopsis thaliana* locus with GenBank accession number AC004557.

20 SEQ ID NO 15: Amino acid sequence of the *Arabidopsis thaliana* putative open reading frame AAD10668 derived from the *Arabidopsis thaliana* locus with GenBank accession number AC003027.

Further provisional and/or partial sequences are defined through the following SEQ IDs:

25 SEQ ID NO. 1a: The amino acid sequence of the yeast ORF YNR008w from *Saccharomyces cerevisiae*.

SEQ ID NO. 2a: Amino acid sequence of the region of the *Arabidopsis thaliana* genomic sequence (AC004557).

30

SEQ ID NO. 3a: Amino acid sequence of the region of the *Arabidopsis thaliana* genomic sequence (AB006704).

5 SEQ ID NO. 4a: The corresponding genomic DNA sequence and amino acid sequence of the yeast ORF YNROO8w from *Saccharomyces cerevisiae*.

10 SEQ ID NO. 5a: The amino acid sequence of the yeast ORF YNROO8w from *Saccharomyces cerevisiae* derived from the corresponding genomic DNA sequence.

15 SEQ ID NO. 1b: Genomic DNA sequence of the *Saccharomyces cerevisiae* PDAT gene, YNR008w, genebank nucleotide ID number 1302481, and the suggested YNR008w amino acid sequence.

SEQ ID NO. 2b: The suggested amino acid sequence of the yeast gene YNR008w from *Saccharomyces cerevisiae*.

20 SEQ ID NO. 3b: Genomic DNA sequence of the *Schizosaccharomyces pombe* gene SPBC776.14.

SEQ ID NO. 4b: Genomic DNA sequence of part of the *Arabidopsis thaliana* locus with genebank accession number AB006704.

25 SEQ ID NO. 5b: Nucleotide sequence and the corresponding amino acid sequence of the *Arabidopsis thaliana* EST-clone with genebank accession number T04806, and ID number 315966.

30 SEQ ID NO. 6b: Nucleotide and amino acid sequence of the *Zea mays* cDNA clone with genebank ID number 4388167.

SEQ ID NO. 7b: Amino acid sequence of the *Zea mays* cDNA clone with genebank ID number 4388167.

5 SEQ ID NO. 8b: DNA sequence of part of the *Neurospora crassa* cDNA clone WO7G1, ID number 4241729.

SEQ ID NO. 9b: Genomic DNA sequence of part of the *Arabidopsis thaliana* locus with genebank accession number AC004557.

10 SEQ ID NO. 10b: Genomic DNA sequence of part of the *Arabidopsis thaliana* locus with genebank accession number AC003027.

SEQ ID NO. 11b: DNA sequence of part of the *Lycopersicon esculentum* cDNA clone with genebank accession number AI486635.  
15

20

25

30

**Claims**

1. An enzyme catalysing in an acyl-CoA-independent reaction the transfer of fatty acids from phospholipids to diacylglycerol in the biosynthetic pathway for the production of triacylglycerol.
2. An enzyme according to claim 1, comprising an amino acid sequence as set forth in SEQ ID No. 2 or a functional fragment, derivate, allele, homolog or isoenzyme thereof.
3. An enzyme according to claims 1 or 2 designated as phospholipid:diacylglycerol acyltransferase (PDAT).
4. An enzyme according to claims 1 to 3, comprising an amino acid sequence as set forth in SEQ ID No. 1a, 2b or 5a or a functional fragment, derivate, allele, homolog or isoenzyme thereof.
5. An enzyme according to claims 1 to 4, comprising an amino acid sequence selected from the group consisting of sequences as set forth in SEQ ID No. 2a, 3a, 5b, 6, 7b, 8, 13, 14 or 15 or a functional fragment, derivate, allele, homolog or isoenzyme thereof.
6. An enzyme according to claims 1 to 5, comprising an amino acid sequence encoded through a nucleotide sequence, a portion, derivate, allele or homolog thereof selected from the group consisting of sequences as set forth in SEQ ID No. 1, 1b, 3, 3b, 4, 4a, 4b, 5, 5b, 6b, 7, 8b, 9, 9b, 10, 10b, 11, 11b or 12 or a functional fragment, derivate, allele, homolog or isoenzyme of the enzyme encoding amino acid sequence.
7. A nucleotide sequence encoding an enzyme catalysing in an acyl-CoA-independent reaction the transfer of fatty acids from phospholipids to

diacylglycerol in the biosynthetic pathway for the production of triacylglycerol.

- 5 8. A nucleotide sequence according to claim 7 encoding an enzyme designated as phospholipid:diacylglycerol acyltransferase (PDAT).
9. A nucleotide sequence according to claims 7 or 8, selected from the group consisting of sequences as set forth in SEQ ID No. 1, 1b, 3, 3b, 4, 4a, 4b, 9b, 10, 10b or 11 or a portion, derivate, allele or homolog thereof.
- 10 10. A partial nucleotide sequence corresponding to a fulllength nucleotide sequence according to claims 7 to 9, selected from the group consisting of sequences as set forth in SEQ ID No. 5, 5b, 6b, 7, 8b, 9, 11b or 12 or a portion, derivate, allele or homolog thereof.
- 15 11. A nucleotide sequence according to claims 7 to 10, comprising a nucleotide sequence which is at least 40% homologous to a nucleotide sequence selected form the group consisting of those sequences set forth in SEQ ID No. 1, 1b, 3, 3b, 4, 4a, 4b, 5, 5b, 6b, 7, 8b, 9, 9b, 10, 10b, 11, 11b or 12.
- 20 12. A gene construct comprising a nucleotide sequence according to claims 7 to 11 operably linked to a heterologous nucleic acid.
13. A vector comprising a nucleotide sequence according to claims 7 to 11 or a gene construct according to claim 12.
- 25 14. A vector according to claim 13, which is an expression vector.
15. A vector according to claims 13 or 14, further comprising a selectable marker gene and/or nucleotide sequences for the replication in a host cell or the integration into the genome of the host cell.
- 30

16. A transgenic cell or organism containing a nucleotide sequence according to claims 7 to 11 and/or a gene construct according to claim 12 and/or a vector according to claims 13 to 15.
- 5 17. A transgenic cell or organism according to claim 16 which is an eucaryotic cell or organism.
18. A transgenic cell or organism according to claims 16 or 17 which is a yeast  
10 cell or a plant cell or a plant.
19. A transgenic cell or organism according to claims 16 to 18 having an altered biosynthetic pathway for the production of triacylglycerol.
- 15 20. A transgenic cell or organism according to claims 16 to 19 having an altered oil content.
21. A transgenic cell or organism according to claims 16 to 20 wherein the activity of PDAT is altered.
- 20 22. A transgenic cell or organism according to claims 16 to 21 wherein the altered activity of PDAT is characterized by an alteration in gene expression, catalytic activity and/or regulation of activity of the enzyme.
- 25 23. A transgenic cell or organism according to claims 16 to 22 wherein the altered biosynthetic pathway for the production of triacylglycerol is characterized by the prevention of accumulation of undesirable fatty acids in the membrane lipids.
- 30 24. A process for the production of triacylglycerol, comprising growing a transgenic cell or organism according to claims 16 to 23 under conditions

whereby the said nucleotide sequence according to claims 7 to 11 is expressed.

25. Triacylglycerols produced by a process according to claim 24.

5

26. Use of a nucleotide sequence according to claims 7 to 11 and/or an enzyme according to claims 1 to 6 for the production of triacylglycerol and/or triacylglycerols with uncommon fatty acids.


10 27. Use of a nucleotide sequence according to claims 7 to 11 and/or an enzyme according to claims 1 to 6 for the transformation of any cell or organism in order to be expressed in this cell or organism and result in an altered, preferably increased oil content of this cell or organism.


15








Figures

Fig. 1:



**Fig 2**





## Tables

Tab. 1:

Substrate added	[ <sup>14</sup> C]-lipid <sup>(2)</sup>	unlabelled lipid <sup>(2)</sup>	mol % of added [ <sup>14</sup> C]-acyl group in TAG <sup>(1)</sup>				
			1-OH-TAG	2-OH-TAG	1-OH-1-ver-TAG	1-OH-2-ver-TAG	3-OH-TAG
A	mono-[ <sup>14</sup> C]-ricinoleoyl-DAG	mono-ricinoleoyl-DAG	2,8	12,4	-	-	-
A	mono-[ <sup>14</sup> C]-ricinoleoyl-DAG	mono-vernoleoyl-DAG	3,2	12,1	1,3	-	-
A	mono-[ <sup>14</sup> C]-ricinoleoyl-DAG	di-vernoleoyl-DAG	4	10	0,5	1,2	-
A	mono-[ <sup>14</sup> C]-ricinoleoyl-DAG	di-ricinoleoyl-PC	0,3	24,8	-	-	-
B	mono-[ <sup>14</sup> C]-ricinoleoyl-PC	none	6,8	8,0	-	-	4,7
C	mono-[ <sup>14</sup> C]-ricinoleoyl-PC	di-oleoyl-DAG	8,6	9,8	-	-	5,0
C	mono-[ <sup>14</sup> C]-ricinoleoyl-PC	mono-ricinoleoyl-DAG	5,7	16,7	-	-	1,9
C	mono-[ <sup>14</sup> C]-ricinoleoyl-PC	di-ricinoleoyl-DAG	4,5	9,4	-	-	9,5
C	mono-[ <sup>14</sup> C]-ricinoleoyl-PC	mono-vernoleoyl-DAG	6,0	11,5	10,9	0,5	7,4
C	mono-[ <sup>14</sup> C]-ricinoleoyl-PC	di-vernoleoyl-DAG	6,7	10,8	1,1	8,4	6,8

Tab. 2:

T1 plant deviation	T2 plant number	nmol fatty acids per mg seed	standard
32-4	1	1277	$\pm 11$ (n=2)
	4	1261	$\pm 63$ (n=3)
	5	1369	$\pm 17$ (n=3)
	6	1312	$\pm 53$ (n=4)
	7	1197	$\pm 54$ (n=5)
	8	1240	$\pm 78$ (n=4)
	9	1283	$\pm 54$ (n=5)
	10	1381	$\pm 35$ (n=5)
26-14	1	1444	$\pm 110$ (n=4)
	2	1617*	$\pm 109$ (n=4)
	3	1374	$\pm 37$ (n=2)
	5	1562*	$\pm 70$ (n=4)
	6	1393	$\pm 77$ (n=4)
	7	1433	$\pm 98$ (n=4)
	8	1581*	$\pm 82$ (n=4)

## Sequence Listing

&lt;210&gt; 1

&lt;211&gt; 1986

&lt;212&gt; genomic DNA

<213> *Saccharomyces cerevisiae*

&lt;221&gt; CDS

&lt;222&gt; (1)..(1983)

&lt;400&gt; 1

```

atg ggc aca ctg ttt cga aga aat gtc cag aac caa aag agt gat tct      48
Met Gly Thr Leu Phe Arg Arg Asn Val Gln Asn Gln Lys Ser Asp Ser
   1             5             10             15

gat gaa aac aat aaa ggg ggt tct gtt cat aac aag cga gag agc aga      96
Asp Glu Asn Asn Lys Gly Gly Ser Val His Asn Lys Arg Glu Ser Arg
           20           25           30

aac cac att cat cat caa cag gga tta ggc cat aag aga aga agg ggt      144
Asn His Ile His His Gln Gln Gly Leu Gly His Lys Arg Arg Arg Gly
       35           40           45

att agt ggc agt gca aaa aga aat gag cgt ggc aaa gat ttc gac agg      192
Ile Ser Gly Ser Ala Lys Arg Asn Glu Arg Gly Lys Asp Phe Asp Arg
       50           55           60

aaa aga gac ggg aac ggt aga aaa cgt tgg aga gat tcc aga aga ctg      240
Lys Arg Asp Gly Asn Gly Arg Lys Arg Trp Arg Asp Ser Arg Arg Leu
       65           70           75           80

att ttc att ctt ggt gca ttc tta ggt gta ctt ttg ccg ttt agc ttt      288
Ile Phe Ile Leu Gly Ala Phe Leu Gly Val Leu Leu Pro Phe Ser Phe
           85           90           95

ggc gct tat cat gtt cat aat agc gat agc gac ttg ttt gac aac ttt      336
Gly Ala Tyr His Val His Asn Ser Asp Ser Asp Leu Phe Asp Asn Phe
       100          105          110

gta aat ttt gat tca ctt aaa gtg tat ttg gat gat tgg aaa gat gtt      384
Val Asn Phe Asp Ser Leu Lys Val Tyr Leu Asp Asp Trp Lys Asp Val
       115          120          125

ctc cca caa ggt ata agt tcg ttt att gat gat att cag gct ggt aac      432
Leu Pro Gln Gly Ile Ser Ser Phe Ile Asp Asp Ile Gln Ala Gly Asn
       130          135          140

```

tac tcc aca tct tct tta gat gat ctc agt gaa aat ttt gcc gtt ggt	480
Tyr Ser Thr Ser Ser Leu Asp Asp Leu Ser Glu Asn Phe Ala Val Gly	
145 150 155 160	
aaa caa ctc tta cgt gat tat aat atc gag gcc aaa cat cct gtt gta	528
Lys Gln Leu Leu Arg Asp Tyr Asn Ile Glu Ala Lys His Pro Val Val	
165 170 175	
atg gtt cct ggt gtc att tct acg gga att gaa agc tgg gga gtt att	576
Met Val Pro Gly Val Ile Ser Thr Gly Ile Glu Ser Trp Gly Val Ile	
180 185 190	
gga gac gat gag tgc gat agt tct gcg cat ttt cgt aaa cgg ctg tgg	624
Gly Asp Asp Glu Cys Asp Ser Ser Ala His Phe Arg Lys Arg Leu Trp	
195 200 205	
gga agt ttt tac atg ctg aga aca atg gtt atg gat aaa gtt tgt tgg	672
Gly Ser Phe Tyr Met Leu Arg Thr Met Val Met Asp Lys Val Cys Trp	
210 215 220	
ttg aaa cat gta atg tta gat cct gaa aca ggt ctg gac cca ccg aac	720
Leu Lys His Val Met Leu Asp Pro Glu Thr Gly Leu Asp Pro Pro Asn	
225 230 235 240	
ttt acg cta cgt gca gca cag ggc ttc gaa tca act gat tat ttc atc	768
Phe Thr Leu Arg Ala Ala Gln Gly Phe Glu Ser Thr Asp Tyr Phe Ile	
245 250 255	
gca ggg tat tgg att tgg aac aaa gtt ttc caa aat ctg gga gta att	816
Ala Gly Tyr Trp Ile Trp Asn Lys Val Phe Gln Asn Leu Gly Val Ile	
260 265 270	
ggc tat gaa ccc aat aaa atg acg agt gct gcg tat gat tgg agg ctt	864
Gly Tyr Glu Pro Asn Lys Met Thr Ser Ala Ala Tyr Asp Trp Arg Leu	
275 280 285	
gca tat tta gat cta gaa aga cgc gat agg tac ttt acg aag cta aag	912
Ala Tyr Leu Asp Leu Glu Arg Arg Asp Arg Tyr Phe Thr Lys Leu Lys	
290 295 300	
gaa caa atc gaa ctg ttt cat caa ttg agt ggt gaa aaa gtt tgt tta	960
Glu Gln Ile Glu Leu Phe His Gln Leu Ser Gly Glu Lys Val Cys Leu	
305 310 315 320	
att gga cat tct atg ggt tct cag att atc ttt tac ttt atg aaa tgg	1008
Ile Gly His Ser Met Gly Ser Gln Ile Ile Phe Tyr Phe Met Lys Trp	
325 330 335	



gtc gag gct gaa ggc cct ctt tac ggt aat ggt ggt cgt ggc tgg gtt	1056
Val Glu Ala Glu Gly Pro Leu Tyr Gly Asn Gly Gly Arg Gly Trp Val	
340 345 350	
aac gaa cac ata gat tca ttc att aat gca gca ggg acg ctt ctg ggc	1104
Asn Glu His Ile Asp Ser Phe Ile Asn Ala Ala Gly Thr Leu Leu Gly	
355 360 365	
gct cca aag gca gtt cca gct cta att agt ggt gaa atg aaa gat acc	1152
Ala Pro Lys Ala Val Pro Ala Leu Ile Ser Gly Glu Met Lys Asp Thr	
370 375 380	
att caa tta aat acg tta gcc atg tat ggt ttg gaa aag ttc ttc tca	1200
Ile Gln Leu Asn Thr Leu Ala Met Tyr Gly Leu Glu Lys Phe Phe Ser	
385 390 395 400	
aga att gag aga gta aaa atg tta caa acg tgg ggt ggt ata cca tca	1248
Arg Ile Glu Arg Val Lys Met Leu Gln Thr Trp Gly Gly Ile Pro Ser	
405 410 415	
atg cta cca aag gga gaa gag gtc att tgg ggg gat atg aag tca tct	1296
Met Leu Pro Lys Gly Glu Glu Val Ile Trp Gly Asp Met Lys Ser Ser	
420 425 430	
tca gag gat gca ttg aat aac aac act gac aca tac ggc aat ttc att	1344
Ser Glu Asp Ala Leu Asn Asn Asn Thr Asp Thr Tyr Gly Asn Phe Ile	
435 440 445	
cga ttt gaa agg aat acg agc gat gct ttc aac aaa aat ttg aca atg	1392
Arg Phe Glu Arg Asn Thr Ser Asp Ala Phe Asn Lys Asn Leu Thr Met	
450 455 460	
aaa gac gcc att aac atg aca tta tcg ata tca cct gaa tgg ctc caa	1440
Lys Asp Ala Ile Asn Met Thr Leu Ser Ile Ser Pro Glu Trp Leu Gln	
465 470 475 480	
aga aga gta cat gag cag tac tcg ttc ggc tat tcc aag aat gaa gaa	1488
Arg Arg Val His Glu Gln Tyr Ser Phe Gly Tyr Ser Lys Asn Glu Glu	
485 490 495	
gag tta aga aaa aat gag cta cac cac aag cac tgg tcg aat cca atg	1536
Glu Leu Arg Lys Asn Glu Leu His His Lys His Trp Ser Asn Pro Met	
500 505 510	
gaa gta cca ctt cca gaa gct ccc cac atg aaa atc tat tgt ata tac	1584
Glu Val Pro Leu Pro Glu Ala Pro His Met Lys Ile Tyr Cys Ile Tyr	
515 520 525	

ggg gtg aac aac cca act gaa agg gca tat gta tat aag gaa gag gat	1632
Gly Val Asn Asn Pro Thr Glu Arg Ala Tyr Val Tyr Lys Glu Glu Asp	
530 535 540	
 gac tcc tct gct ctg aat ttg acc atc gac tac gaa agc aag caa cct	1680
Asp Ser Ser Ala Leu Asn Leu Thr Ile Asp Tyr Glu Ser Lys Gln Pro	
545 550 555 560	
 gta ttc ctc acc gag ggg gac gga acc gtt ccg ctc gtg gcg cat tca	1728
Val Phe Leu Thr Glu Gly Asp Gly Thr Val Pro Leu Val Ala His Ser	
565 570 575	
 atg tgt cac aaa tgg gcc cag ggt gct tca ccg tac aac cct gcc gga	1776
Met Cys His Lys Trp Ala Gln Gly Ala Ser Pro Tyr Asn Pro Ala Gly	
580 585 590	
 att aac gtt act att gtg gaa atg aaa cac cag cca gat cga ttt gat	1824
Ile Asn Val Thr Ile Val Glu Met Lys His Gln Pro Asp Arg Phe Asp	
595 600 605	
 ata cgt ggt gga gca aaa agc gcc gaa cac gta gac atc ctc ggc agc	1872
Ile Arg Gly Gly Ala Lys Ser Ala Glu His Val Asp Ile Leu Gly Ser	
610 615 620	
 gcg gag ttg aac gat tac atc ttg aaa att gca agc ggt aat ggc gat	1920
Ala Glu Leu Asn Asp Tyr Ile Leu Lys Ile Ala Ser Gly Asn Gly Asp	
625 630 635 640	
 ctc gtc gag cca cgc caa ttg tct aat ttg agc cag tgg gtt tct cag	1968
Leu Val Glu Pro Arg Gln Leu Ser Asn Leu Ser Gln Trp Val Ser Gln	
645 650 655	
 atg ccc ttc cca atg taa	1986
Met Pro Phe Pro Met	
660	

&lt;210&gt; 2

&lt;211&gt; 661

&lt;212&gt; PRT

<213> *Saccharomyces cerevisiae*

&lt;400&gt; 2

```

Met Gly Thr Leu Phe Arg Arg Asn Val Gln Asn Gln Lys Ser Asp Ser
  1              5              10              15
Asp Glu Asn Asn Lys Gly Gly Ser Val His Asn Lys Arg Glu Ser Arg
      20              25              30
Asn His Ile His His Gln Gln Gly Leu Gly His Lys Arg Arg Arg Gly
      35              40              45
Ile Ser Gly Ser Ala Lys Arg Asn Glu Arg Gly Lys Asp Phe Asp Arg
      50              55              60
Lys Arg Asp Gly Asn Gly Arg Lys Arg Trp Arg Asp Ser Arg Arg Leu
      65              70              75              80
Ile Phe Ile Leu Gly Ala Phe Leu Gly Val Leu Leu Pro Phe Ser Phe
      85              90              95
Gly Ala Tyr His Val His Asn Ser Asp Ser Asp Leu Phe Asp Asn Phe
      100              105              110
Val Asn Phe Asp Ser Leu Lys Val Tyr Leu Asp Asp Trp Lys Asp Val
      115              120              125
Leu Pro Gln Gly Ile Ser Ser Phe Ile Asp Asp Ile Gln Ala Gly Asn
      130              135              140
Tyr Ser Thr Ser Ser Leu Asp Asp Leu Ser Glu Asn Phe Ala Val Gly
      145              150              155              160
Lys Gln Leu Leu Arg Asp Tyr Asn Ile Glu Ala Lys His Pro Val Val
      165              170              175
Met Val Pro Gly Val Ile Ser Thr Gly Ile Glu Ser Trp Gly Val Ile
      180              185              190
Gly Asp Asp Glu Cys Asp Ser Ser Ala His Phe Arg Lys Arg Leu Trp
      195              200              205
Gly Ser Phe Tyr Met Leu Arg Thr Met Val Met Asp Lys Val Cys Trp
      210              215              220
Leu Lys His Val Met Leu Asp Pro Glu Thr Gly Leu Asp Pro Pro Asn
      225              230              235              240
Phe Thr Leu Arg Ala Ala Gln Gly Phe Glu Ser Thr Asp Tyr Phe Ile
      245              250              255
Ala Gly Tyr Trp Ile Trp Asn Lys Val Phe Gln Asn Leu Gly Val Ile
      260              265              270
Gly Tyr Glu Pro Asn Lys Met Thr Ser Ala Ala Tyr Asp Trp Arg Leu
      275              280              285
Ala Tyr Leu Asp Leu Glu Arg Arg Asp Arg Tyr Phe Thr Lys Leu Lys
      290              295              300
Glu Gln Ile Glu Leu Phe His Gln Leu Ser Gly Glu Lys Val Cys Leu
      305              310              315              320
Ile Gly His Ser Met Gly Ser Gln Ile Ile Phe Tyr Phe Met Lys Trp
      325              330              335

```

SUBSTITUTE SHEET (RULE 26)

Val Glu Ala Glu Gly Pro Leu Tyr Gly Asn Gly Gly Arg Gly Trp Val  
 340 345 350  
 Asn Glu His Ile Asp Ser Phe Ile Asn Ala Ala Gly Thr Leu Leu Gly  
 355 360 365  
 Ala Pro Lys Ala Val Pro Ala Leu Ile Ser Gly Glu Met Lys Asp Thr  
 370 375 380  
 Ile Gln Leu Asn Thr Leu Ala Met Tyr Gly Leu Glu Lys Phe Phe Ser  
 385 390 395 400  
 Arg Ile Glu Arg Val Lys Met Leu Gln Thr Trp Gly Gly Ile Pro Ser  
 405 410 415  
 Met Leu Pro Lys Gly Glu Glu Val Ile Trp Gly Asp Met Lys Ser Ser  
 420 425 430  
 Ser Glu Asp Ala Leu Asn Asn Asn Thr Asp Thr Tyr Gly Asn Phe Ile  
 435 440 445  
 Arg Phe Glu Arg Asn Thr Ser Asp Ala Phe Asn Lys Asn Leu Thr Met  
 450 455 460  
 Lys Asp Ala Ile Asn Met Thr Leu Ser Ile Ser Pro Glu Trp Leu Gln  
 465 470 475 480  
 Arg Arg Val His Glu Gln Tyr Ser Phe Gly Tyr Ser Lys Asn Glu Glu  
 485 490 495  
 Glu Leu Arg Lys Asn Glu Leu His His Lys His Trp Ser Asn Pro Met  
 500 505 510  
 Glu Val Pro Leu Pro Glu Ala Pro His Met Lys Ile Tyr Cys Ile Tyr  
 515 520 525  
 Gly Val Asn Asn Pro Thr Glu Arg Ala Tyr Val Tyr Lys Glu Glu Asp  
 530 535 540  
 Asp Ser Ser Ala Leu Asn Leu Thr Ile Asp Tyr Glu Ser Lys Gln Pro  
 545 550 555 560  
 Val Phe Leu Thr Glu Gly Asp Gly Thr Val Pro Leu Val Ala His Ser  
 565 570 575  
 Met Cys His Lys Trp Ala Gln Gly Ala Ser Pro Tyr Asn Pro Ala Gly  
 580 585 590  
 Ile Asn Val Thr Ile Val Glu Met Lys His Gln Pro Asp Arg Phe Asp  
 595 600 605  
 Ile Arg Gly Gly Ala Lys Ser Ala Glu His Val Asp Ile Leu Gly Ser  
 610 615 620  
 Ala Glu Leu Asn Asp Tyr Ile Leu Lys Ile Ala Ser Gly Asn Gly Asp  
 625 630 635 640  
 Leu Val Glu Pro Arg Gln Leu Ser Asn Leu Ser Gln Trp Val Ser Gln  
 645 650 655  
 Met Pro Phe Pro Met  
 660

<210> 3  
<211> 2312  
<212> genomic DNA  
<213> Schizosaccharomyces pombe

<400> 3  
ATGGCGTCTT CCAAGAAGAG CAAAACATCAT AAGAAAAAGA AAGAAGTCAA 50  
ATCTCCTATC GACTTACCAA ATTCAAAGAA ACCAACTCGC GCTTTGAGTG 100  
AGCAACCTTC AGCGTCCGAA ACACAATCTG TTTCAAATAA ATCAAGAAAA 150  
TCTAAATTTG GAAAAAGATT GAATTTTATA TTGGGCGCTA TTTTGGAAT 200  
ATGCGGTGCT TTTTTTTTCG CTGTTGGAGA CGACAATGCT GTTTTCGACC 250  
CTGCTACGTT AGATAAATTT GGAATATGCT TAGGCTCTTC AGACTTGTTT 300  
GATGACATTA AAGGATATTT ATCTTATAAT GTGTTTAAGG ATGCACCTTT 350  
TACTACGGAC AAGCCTTCGC AGTCTCCTAG CGGAAATGAA GTTCAAGTTG 400  
GTCTTGATAT GTACAATGAG GGATATCGAA GTGACCATCC TGTTATTATG 450  
GTTCTGGTG TTATCAGCTC AGGATTAGAA AGTTGGTCGT TTAATAATTG 500  
CTCGATTCTT TACTTTAGGA AACGTCTTTG GGGTAGCTGG TCTATGCTGA 550  
AGGCAATGTT CCTTGACAAG CAATGCTGGC TTGAACATTT AATGCTTGAT 600  
AAAAAAACCG GCTTGGATCC GAAGGGAATT AAGCTGCGAG CAGCTCAGGG 650  
GTTTGAAGCA GCTGATTTTT TTATCACGGG CTATTGGATT TGGAGTAAAG 700  
TAATTGAAAA CCTTGCTGCA ATTGGTTATG AGCCTAATAA CATGTTAAGT 750  
GCTTCTTACG ATTGGCGGTT ATCATATGCA AATTTAGAGG AACGTGATAA 800  
ATATTTTTCA AAGTTAAAAA TGTTCAATTGA GTACAGCAAC ATTGTACATA 850  
AGAAAAAGGT AGTGTTGATT TCTCACTCCA TGGGTTTACA GGTTACGTAC 900  
TATTTTTTTA AGTGGGTTGA AGCTGAGGGC TACGGAAATG GTGGACCGAC 950  
TTGGGTTAAT GATCATATTG AAGCATTATG AAATGTGAGT CTCGATGGTT 1000  
GTTTGACTAC GTTTCTAACT TTTGAATAGA TATCGGGATC TTTGATTGGA 1050  
GCACCCAAAA CAGTGGCAGC GCTTTTATCG GGTGAAATGA AAGATACAGG 1100  
TATTGTAATT ACATTAAACA TGTTAATATT TAATTTTTGC TAACCGTTTT 1150  
AAGCTCAATT GAATCAGTTT TCGGTCTATG GGTAAAGCAAT AAATTGTTGA 1200  
GATTTGTTAC TAATTTACTG TTTAGTTTGG AAAAATTTTT TTCCCGTTCT 1250  
GAGGTATATT CAAAAATACA AATGTGCTCT ACTTTTTCTA ACTTTTAATA 1300  
GAGAGCCATG ATGGTTCGCA CTATGGGAGG AGTTAGTTCT ATGCTTCCTA 1350  
AAGGAGGCGA TGTTGTATGG GGAAATGCCA GTTGGGTAAG AAATATGTGC 1400  
TGTTAATTTT TTATTAATAT TTAGGCTCCA GATGATCTTA ATCAAACAAA 1450  
TTTTTCCAAT GGTGCAATTA TTCGATATAG AGAAGACATT GATAAGGACC 1500  
ACGATGAATT TGACATAGAT GATGCATTAC AATTTTTTAA AAATGTTACA 1550  
GATGACGATT TTAAAGTCAT GCTAGCGAAA AATTATTCCC ACGGTCTTGC 1600  
TTGGACTGAA AAAGAAGTGT TAAAAAATAA CGAAATGCCG TCTAAATGGA 1650  
TAAATCCGCT AGAAGTAAGA ACATTAAAGT TACTAAATTA TACTAACCCA 1700  
AATAGACTAG TCTTCCTTAT GCTCCTGATA TGAAAATTTA TTGCGTTTAC 1750  
GGGGTCGGAA AACCAACTGA GAGAGGTTAT TATTATACTA ATAATCCTGA 1800  
GGGGCAACCT GTCATTGATT CCTCGGTTAA TGATGGAACA AAAGTTGAAA 1850  
ATGTGAGAGA ATTTATGTTT CAAACATTCT ATTAAGTGT TTATTAGGGT 1900  
ATTGTTATGG ATGATGGTGA TGGAACCTTA CCAATATTAG CCCTTGTTT 1950  
GGTGTGCAAT AAAGTTTGGC AAACAAAAAG GTTTAATCCT GCTAATACAA 2000  
GTATCACAAA TTATGAAATC AAGCATGAAC CTGCTGCGTT TGATCTGAGA 2050  
GGAGGACCTC GCTCGGCAGA ACACGTCGAT ATACTTGAC ATTACAGAGCT 2100  
AAATGTATGT TCATTTTACC TTACAAATTT CTATTACTAA CTCTTGAAAT 2150  
AAGGAAATTA TTTTAAAAGT TTCATCAGGC CATGGTGACT CGGTACCAAA 2200  
CCGTTATATA TCAGATATCC AGTACGGACA TAAGTTTGT AGATTGCAAT 2250  
TAACTAATA ACCGAACAGG GAAATAATAA ATGAGATAAA TCTCGATAAA 2300  
CCTAGAAATT AA 2312

<210> 4  
 <211> 3685  
 <212> genomic DNA  
 <213> Arabidopsis thaliana

<400> 4

ATGCCCCCTTA	TTCATCGGAA	AAAGCCGACG	GAGAAACCAT	CGACGCCGCC	50
ATCTGAAGAG	GTGGTGCACG	ATGAGGATTC	GCAAAAAGAAA	CCACACGAAT	100
CTTCCAAATC	CCACCATAAG	AAATCGAACG	GAGGAGGGAA	GTGGTCGTGC	150
ATCGATTCTT	GTTGTTGGTT	CATTGGGTGT	GTGTGTGTAA	CCTGGTGGTT	200
TCTTCTCTTC	CTTTACAACG	CAATGCCTGC	GAGCTTCCCT	CAGTATGTAA	250
CGGAGCGAAT	CACGGGTCCT	TTGCCTGACC	CGCCCCGGTG	TAAGCTCAAA	300
AAAGAAGGTC	TTAAGGCGAA	ACATCCTGTT	GTCTTCATTC	CTGGGATTGT	350
CACCGGTGGG	CTCGAGCTTT	GGGAAGGCAA	ACAATGCGCT	GATGGTTTAT	400
TTAGAAAACG	TTTGTGGGGT	GGAACCTTTG	GTGAAGTCTA	CAAAAGGTGA	450
GCTCAACAAT	TCTCACTCTT	CCTTTATATT	GGGATTTGGA	TTGGATCTGA	500
TGAGATCACG	CACCTGTTGC	TTCTTCAACA	TCACTCAAAC	TTTAATTCCA	550
TGTTTGTCTG	TCTTACTCTT	TACTTTTTTT	TTTTTTTGAT	GTGAAACGCT	600
ATTTTCTTAA	GAGACTATTT	CTGTATGTGT	AAGGTAAGCG	TTCCAAGGAC	650
GTAATTGGCT	TGGACTATTT	CTGTTTGATT	GTTAACTTTA	GGATATAAAA	700
TAGCTGCCTT	GGAATTTCAA	GTCATCTTAT	TGCCAAATCT	GTTGCTAGAC	750
ATGCCCTAGA	GTCCGTTTCAT	AACAAGTTAC	TTCCTTTACT	GTCGTTGCGT	800
GTAGATTTAG	CTTTGTGTAG	CGTATAATGA	AGTAGTGTTT	TATGTTTGTG	850
TGGGAATAGA	GAAGTTCTAA	CTACATCTGT	GGAAAGTGTG	TTCAGGCTGT	900
GATAGAGGAC	TGTTGCTTTA	TTATTCAACT	ATGTATATGT	GTAATTAAAG	950
CTAGTTCCTT	TTTGATCTTT	CAGCTCAATG	TGCTTTTCTC	AATTTTTTTC	1000
TCAATTTCAA	AGTTTCACAT	CGAGTTTAT	CACATGTCTT	GAATTTTCGT	1050
CATCCTCGTT	CTGTTATCCA	GCTTTGAACT	CCTCCCGACC	CTGCTATGGA	1100
TATATTAAAA	AAAAAGTGTT	TTGTGGGTTG	CATCTTTGTT	ACGATCTGCA	1150
TCTTCTTCTT	TCCGCTCAGT	GTTTCATGTT	TTGCTATGGT	AGAGATGGGC	1200
AATGTTATTG	TTGATGGTAA	CAGTGGTATA	GTTGATAGTA	TCTTAACTAA	1250
TCAATTATCT	CTTTGATTCA	GGCCTCTATG	TTGGGTGGAA	CACATGTCAC	1300
TTGACAATGA	AACTGGGTTG	GATCCAGCTG	GTATTAGAGT	TCGAGCTGTA	1350
TCAGGACTCG	TGGTGTCTGA	CTACTTTGCT	CCTGGCTACT	TTGTCTGGGC	1400
AGTGCTGATT	GCTAACCTTG	CACATATTGG	ATATGAAGAG	AAAAATATGT	1450
ACATGGCTGC	ATATGACTGG	CGGCTTTCGT	TTCAGAACAC	AGAGGTTCTT	1500
TTCTCATCGT	TCTTTCTATT	ATTCTGTTCC	ATGTTACGTT	TCTTTCTTCA	1550
TTACTTAAGG	CTTAAATATG	TTTCATGTTG	AATTAATAGG	TACGTGATCA	1600
GACTCTTAGC	CGTATGAAAA	GTAATATAGA	GTTGATGGTT	TCTACCAACG	1650
GTGGAAAAAA	AGCAGTTATA	GTTCCGCATT	CCATGGGGGT	CTTGATTTTT	1700
CTACATTTTA	TGAAGTGGGT	TGAGGCACCA	GCTCCTCTGG	GTGGCGGGGG	1750
TGGGCCAGAT	TGGTGTGCAA	AGTATATTAA	GGCGGTGATG	AACATTGGTG	1800
GACCATTTCT	TGGTGTTCCT	AAAGCTGTTG	CAGGGCTTTT	CTCTGCTGAA	1850
GCAAAGGATG	TTGCAGTTGC	CAGGTATTGA	ATATCTGCTT	ATACTTTTGA	1900
TGATCAGAAC	CTTGGCTCTG	GAACTCAAAG	TTATTCTACT	AAATATCAAT	1950
TCTAATAACA	TTGCTATATT	ATCGCTGCAA	CTGACATTGG	TTGATTATTT	2000
TTGCTGCTTA	TGTAAGTCAA	ACTCTCTTGA	GATTAGACAA	ATGATGAATT	2050
GATAATTCTT	ACGCATTGCT	CTGTGATGAC	CAGTTTCTTA	GCTTCGACGA	2100
TAACATTTGT	CATACTGTCT	TTTGGAGGGC	ATTGAATTTT	GCTATGGAAA	2150
GCGCTGGAGC	TTCCATGCTT	GCATTCTTTA	CCAATTAGCG	TTATTCTGCT	2200
TCTTTCAATT	TTCTTGTATA	TGCATCTATG	GTCTTTTATT	TCTTCTTAAT	2250
TAAAGACTCG	TTGGATTAGT	TGCTCTATTA	GTCACTTGGT	TCCTTAATAT	2300
AGAACTTTAC	TTTCTTCGAA	AATTGCAGAG	CGATTGCCCC	AGGATTCTTA	2350
GACACCGATA	TATTTAGACT	TCAGACCTTG	CAGCATGTAA	TGAGAATGAC	2400
ACGCACATGG	GACTCAACAA	TGTCTATGTT	ACCGAAGGGA	GGTGACACGA	2450
TATGGGGCGG	GCTTGATTGG	TCACCGGAGA	AAGGCCACAC	CTGTTGTGGG	2500
AAAAAGCAAA	AGAACAACGA	AACTTGTGGT	GAAGCAGGTG	AAAACGGAGT	2550
TTCCAAGAAA	AGTCCTGTTA	ACTATGGAAG	GATGATATCT	TTTGGGAAAG	2600
AAGTAGCAGA	GGCTGCGCCA	TCTGAGATTA	ATAATATTGA	TTTTCGAGTA	2650
AGGACATATA	AATCATAATA	AACCTTGTAC	ATTTTGTGAT	TGTATGATGA	2700
ATATCTGTAC	ATTTTATCTG	GTGAAGGGTG	CTGTCAAAGG	TCAGAGTATC	2750
CCAAATCACA	CCTGTCTGTA	CGTGTGGACA	GAGTACCATG	ACATGGGAAT	2800
TGCTGGGATC	AAAGCTATCG	CTGAGTATAA	GGTCTACACT	GCTGGTGAAG	2850

CTATAGATCT	ACTACATTAT	GTTGCTCCTA	AGATGATGGC	GCGTGGTGCC	2900
GCTCATTCT	CTTATGGAAT	TGCTGATGAT	TTGGATGACA	CCAAGTATCA	2950
AGATCCCAA	TACTGGTCAA	ATCCGTTAGA	GACAAAGTAA	GTGATTTCTT	3000
GATTCCAAC	GTATCCTTCG	TCCTGATGCA	TTATCAGTCT	TTTTGTTTTT	3050
GGTCTTGTT	GATATGGTTT	TCAGCTCAAA	GCTTACAAAG	CTGTTTCTGA	3100
GCCTTTCTC	AAAAGGCTTG	CTCAGTAATA	TTGAGGTGCT	AAAGTTGATA	3150
CATGTGACT	TTGCTTATAA	ATCCTCCGTT	TGGTTTGTTT	TGCTTTTTC	3200
GATTACCGA	TGCTCCTGAG	ATGGAAATCT	ACTCATTATA	CGGAGTGGGG	3250
ATACCAACG	AACGAGCATA	CGTATACAAG	CTTAACCAGT	CTCCCGACAG	3300
TTGCATCCCC	TTTCAGATAT	TCACTTCTGC	TCACGAGGAG	GACGAAGATA	3350
GCTGTCTGA	AGCAGGAGTT	TACAATGTGG	ATGGGGATGA	AACAGTACCC	3400
GTCCTAAGT	CCGGGTACAT	GTGTGCAAAA	GCGTGGCGTG	GCAAGACAAG	3450
ATTCAACCCT	TCCGGAATCA	AGACTTATAT	AAGAGAATAC	AATCACTCTC	3500
CGCCGGCTAA	CCTGTTGGAA	GGGCGCGGGA	CGCAGAGTGG	TGCCCCATGTT	3550
GATATCATGG	GAAACTTTGC	TTTGATCGAA	GATATCATGA	GGGTTGCCGC	3600
CGGAGGTAAC	GGGTCTGATA	TAGGACATGA	CCAGGTCCAC	TCTGGCATAT	3650
TTGAATGGTC	GGAGCGTATT	GACCTGAAGC	TGTGA		3685

<210> 5  
 <211> 2427  
 <212> cDNA  
 <213> Arabidopsis thaliana

<400> 5  
 AGAAACAGCT CTTTGTCTCT CTCGACTGAT CTAACAATCC CTAATCTGTG 50  
 TTCTAAATTC CTGGACGAGA TTTGACAAAG TCCGTATAGC TTAACCTGGT 100  
 TTAATTTCAA GTGACAGATA TGCCCTTAT TCATCGGAAA AAGCCGACGG 150  
 AGAAACCATC GACGCCGCCA TCTGAAGAGG TGGTGCACGA TGAGGATTCTG 200  
 CAAAAGAAAC CACACGAATC TTCCAAATCC CACCATAAGA AATCGAACGG 250  
 AGGAGGGAAG TGGTCGTGCA TCGATTCTTG TTGTTGGTTC ATTGGGTGTG 300  
 TGTGTGTAAC CTGGTGGTTT CTTCTCTTCC TTTACAACGC AATGCCTGCG 350  
 AGCTTCCCTC AGTATGTAAC GGAGCGAATC ACGGGTCCTT TGCCTGACCC 400  
 GCCCGGTGTT AAGCTCAAAA AAAGAAGGTC TTAAGGCGAA ACATCCTGTT 450  
 GTCTTCATCT CTGGGATTGT CACCGGTGGG CTCGAGCTTT GGAAGGCAA 500  
 ACAATGCGCT GATGGTTTAT TTAGAAAACG TTTGTGGGGT GGAACCTTTTG 550  
 GTGAAGTCTA CAAAAGGCCT CTATGTTGGG TGGAACACAT GTCACCTGAC 600  
 AATGAACTG GGTGATCC AGCTGGTATT AGAGTTCGAG CTGTATCAGG 650  
 ACTCGTGGCT GCTGACTACT TTGCTCCTGG CTACTTTGTC TGGGCAGTGC 700  
 TGATTGCTAA CCTTGCACAT ATTGGATATG AAGAGAAAAA TATGTACATG 750  
 GCTGCATATG ACTGGCGGCT TTCGTTTCAG AACACAGAGG TACGTGATCA 800  
 GACTCTTAGC CGTATGAAAA GTAATATAGA GTTGATGGTT TCTACCAACG 850  
 GTGGAAAAAA AGCAGTTATA GTTCCGCATT CCATGGGGGT CTTGTATTTT 900  
 CTACATTTTA TGAAGTGGGT TGAGGCACCA GCTCCTCTGG GTGGCGGGG 950  
 TGGGCCAGAT TGGTGTGCAA AGTATATTAA GGCGGTGATG AACATTGGTG 1000  
 GACCATTTCT TGGTGTTCOA AAAGCTGTTG CAGGGCTTTT CTCTGCTGAA 1050  
 GCAAAGGATG TTGCAGTTGC CAGAGCGATT GCCCCAGGAT TCTTAGACAC 1100  
 CGATATATTT AGACTTCAGA CCTTGCAGCA TGTAATGAGA ATGACACGCA 1150  
 CATGGGACTC AACAATGTCT ATGTTACCGA AGGGAGGTGA CACGATATGG 1200  
 GGCGGGCTTG ATTGGTCACC GGAGAAAAGC CACACCTGTT GTGGGAAAAA 1250  
 GCAAAAGAAC AACGAAACTT GTGGTGAAGC AGGTGAAAAC GGAGTTTCCA 1300  
 AGAAAAGTCC TGTTAACTAT GGAAGGATGA TATCTTTTGG GAAAGAAGTA 1350  
 GCAGAGGCTG CGCCATCTGA GATTAATAAT ATTGATTTTC GAGGTGCTGT 1400  
 CAAAGGTCAG AGTATCCCAA ATCACACCTG TCGTGACGTG TGGACAGAGT 1450  
 ACCATGACAT GGGAAATTGCT GGGATCAAAG CTATCGCTGA GTATAAGGTC 1500  
 TACACTGCTG GTGAAGCTAT AGATCTACTA CATTATGTTG CTCCTAAGAT 1550  
 GATGGCGCGT GGTGCCGCTC ATTTCTCTTA TGGAATTGCT GATGATTTGG 1600  
 ATGACACCAA GTATCAAGAT CCCAAATACT GGTCAAATCC GTTAGAGACA 1650  
 AAATTACCGA ATGCTCCTGA GATGGAAATC TACTCATTAT ACGGAGTGGG 1700  
 GATACCAACG GAACGAGCAT ACGTATACAA GCTTAACCAG TCTCCCGACA 1750  
 GTTGCATCCC CTTTCAGATA TTCACTTCTG CTCACGAGGA GGACGAAGAT 1800  
 AGCTGTCTGA AAGCAGGAGT TTACAATGTG GATGGGGATG AAACAGTACC 1850  
 CGTCCTAAGT GCCGGGTACA TGTGTGCAA AGCGTGCGGT GGCAAGACAA 1900  
 GATTCAACCC TTCCGGAATC AAGACTTATA TAAGAGAATA CAATCACTCT 1950  
 CCGCCGGCTA ACCTGTTGGA AGGGCGCGGG ACGCAGAGTG GTGCCCATGT 2000  
 TGATATCATG GGAAACTTTG CTTTGATCGA AGATATCATG AGGGTTGCCG 2050  
 CCGGAGGTAA CGGGTCTGAT ATAGGACATG ACCAGGTCCA CTCTGGCATA 2100  
 TTTGAATGGT CGGAGCGTAT TGACCTGAAG CTGTGAATAT CATGATCTCT 2150  
 TTAAGCTGTC CTGTCACTT ATGTGAATCC AATACTTTGA AAGAGAGATC 2200  
 ATCATCAATT CATCATCATC GTCATCATCA TGATGCTCAA CTCACAAAGA 2250  
 AGCCTGAGAA TGATACTTTG GTGCGAAATT CTCAATACCT CTTTAATATT 2300  
 CTTATTGAAT GTAAATTATA CAATCCTATC TAATGTTTGA ACGATAACAC 2350  
 AAAACTTGCT GCNGCCATGT TTGTTTGTCT TGTCAAAAGC ATCAATTTGT 2400  
 GGGTTAAAAA AAAAAAAAAA AAAAAA 2427



<210> 6  
<211> 671  
<212> PRT  
<213> Arabidopsis thaliana

<400> 6  
MPLIHRKKPT EKPSTPPSEE VVHDEDSQKK PHESSKSHHK KSNGGGKWSC 50  
IDSCCWFIGC VCVTWWFLLF LYNAMPASFP QYVTERITGP LPDPPGVKLK 100  
KEGLKAKHPV VFIPGIIVTGG LELWEGKQCA DGLFRKRLWG GTFGEVYKRP 150  
LCWVEHMSLD NETGLDPAGI RVRVAVSGLVA ADYFAPGYFV WAVLIANLAH 200  
IGYEEKNMYM AAYDWRLSFQ NTEVRDQTLS RMKSNIELMV STNGGKKAVI 250  
VPHSMGVLYF LHFMKWVEAP APLGGGGGPD WCAKYIKAVM NIGGPFLGVP 300  
KAVAGLFSAE AKDVAVARAI APGFLDTDIF RLQTLQHVMR MTRTW DSTMS 350  
MLPKGGDTIW GGLDWSPEKG HTCCGKKQKN NETCGEAGEN GVSCKSPVNY 400  
GRMISFGKEV AEAAPSEINN IDFRGAVKGQ SIPNHTCRDV WTEYHDMGIA 450  
GIKAIAEYKV YTAGEAIDLL HYVAPKMMAR GAAHFSYGIA DDLDDTKYQD 500  
PKYWSNPLET KLPNAPEMEI YSLYGVGIPT ERAYVYKLNQ SPDSCIPFQI 550  
FTSAHEEDED SCLKAGVYNV DGDETVPVLS AGYMCAKAWR GKTRFNPSGI 600  
KTYIREYNHS PPANLLEGRG TQSGAHVDIM GNFALIEDIM RVAAGNGSD 650  
IGHDQVHSGI FEWSERIDLK L 671

<210> 7  
 <211> 643  
 <212> cDNA  
 <213> Zea mays

<221> CDS  
 <222> (1)..(402)

<400> 6

```

CGG GAG AAA ATA GCT GCT TTG AAG GGG GGT GTT TAC TTA GCC GAT GGT 48
Arg Glu Lys Ile Ala Ala Leu Lys Gly Gly Val Tyr Leu Ala Asp Gly
  1             5             10             15

GAT GAA ACT GTT CCA GTT CTT AGT GCG GGC TAC ATG TGT GCG AAA GGA 96
Asp Glu Thr Val Pro Val Leu Ser Ala Gly Tyr Met Cys Ala Lys Gly
             20             25             30

TGG CGT GGC AAA ACT CGT TTC AGC CCT GCC GGC AGC AAG ACT TAC GTG 144
Trp Arg Gly Lys Thr Arg Phe Ser Pro Ala Gly Ser Lys Thr Tyr Val
             35             40             45

AGA GAA TAC AGC CAT TCG CCA CCC TCT ACT CTC CTG GAA GGC AGG GGC 192
Arg Glu Tyr Ser His Ser Pro Pro Ser Thr Leu Leu Glu Gly Arg Gly
             50             55             60

ACC CAG AGC GGT GCA CAT GTT GAT ATA ATG GGG AAC TTT GCT CTA ATT 240
Thr Gln Ser Gly Ala His Val Asp Ile Met Gly Asn Phe Ala Leu Ile
  65             70             75             80

GAG GAC GTC ATC AGA ATA GCT GCT GGG GCA ACC GGT GAG GAA ATT GGT 288
Glu Asp Val Ile Arg Ile Ala Ala Gly Ala Thr Gly Glu Glu Ile Gly
             85             90             95

GGC GAT CAG GTT TAT TCA GAT ATA TTC AAG TGG TCA GAG AAA ATC AAA 336
Gly Asp Gln Val Tyr Ser Asp Ile Phe Lys Trp Ser Glu Lys Ile Lys
             100            105            110

TTG AAA TTG TAA CCTATGGGAA GTTAAAGAAG TGCCGACCCG TTTATTGCGTTCC 391
Leu Lys Leu
             115

AAAGTGTCTT GCCTGAGTGC AACTCTGGAT TTTGCTTAAA TATTGTAATT TTTCACGC 449
TTCATTGCTC CCTTTGTCAA ATTTACATTT GACAGGACGC CAATGCGATA CGATGTTG 507
TACCGCTATT TTCAGCATTG TATATTAAAC TGTACAGGTG TAAGTTGCAT TTGCCAGC 565
TGAAATTGTG TAGTCGTTTT CTTTACGATT TAATANCAAG TGGCGGAGCA GTGCCCCA 623
AGCNAAAAAA AAAAAAAAAA
             643

```

<210> 8  
<211> 115  
<212> PRT  
<213> Zea mays

<400> 8  
Arg Glu Lys Ile Ala Ala Leu Lys Gly Gly Val Tyr Leu Ala Asp Gly  
1 5 10 15  
Asp Glu Thr Val Pro Val Leu Ser Ala Gly Tyr Met Cys Ala Lys Gly  
20 25 30  
Trp Arg Gly Lys Thr Arg Phe Ser Pro Ala Gly Ser Lys Thr Tyr Val  
35 40 45  
Arg Glu Tyr Ser His Ser Pro Pro Ser Thr Leu Leu Glu Gly Arg Gly  
50 55 60  
Thr Gln Ser Gly Ala His Val Asp Ile Met Gly Asn Phe Ala Leu Ile  
65 70 75 80  
Glu Asp Val Ile Arg Ile Ala Ala Gly Ala Thr Gly Glu Glu Ile Gly  
85 90 95  
Gly Asp Gln Val Tyr Ser Asp Ile Phe Lys Trp Ser Glu Lys Ile Lys  
100 105 110  
Leu Lys Leu  
115

<210> 9  
<211> 616  
<212> cDNA  
<213> Neurospora crassa

<400> 9  
ggtggcgaag acganggcgg aagttggagg ctaacgagaa tgacnctcgg 50  
agatggatct accctctaga gacacgacta ccnttgcacc cagcctcaag 100  
gtntacngtt tntatgggta ggaagccgac ggagcgagcc tacatctatc 150  
tggcgccccga tcccgggacg acaacgcacg tttagatgac gatcgatacg 200  
actttgactn agggggcacat tgaccacggt gtgatttttg gcgaaggcga 250  
tggcacagtg aaccttatga gtttggggta cctgtgcaat aaggggtgga 300  
aaatgaagag atacaatcct gcgggctcaa aaataaccgt ggtcgagatg 350  
ccgcatgaac cagaacggtt caatccgaga ggagggccga atacggcgga 400  
tcacgtggat attctaggaa ggcagaatct aaacgagtac attcttaaag 450  
tggcggcagg tcgaggcgat acaattgagg attttattac tagtaatatt 500  
cttaaatatg tagaaaaggt tgaaatttat gaagagtaat taaatacggc 550  
acataggtta ctcaatagta tgactaatta aaaaaaaatt ttttttctaa 600  
aaaaaaaaaa aaaaaa 616

<210> 10  
<211> 1562  
<212> genomic DNA  
<213> Arabidopsis thaliana

<400> 10  
ATGAAAAAAAA TATCTTCACA TTATTCGGTA GTCATAGCGA TACTCGTTGT 50  
GGTGACGATG ACCTCGATGT GTCAAGCTGT GGGTAGCAAC GTGTACCCTT 100  
TGATTCTGGT TCCAGGAAAC GGAGGTAACC AGCTAGAGGT ACGGCTGGAC 150  
AGAGAATACA AGCCAAGTAG TGTCTGGTGT AGCAGCTGGT TATATCCGAT 200  
TCATAAGAAG AGTGGTGGAT GGTTTAGGCT ATGGTTCGAT GCAGCAGTGT 250  
TATTGTCTCC CTTCACCAGG TGCTTCAGCG ATCGAATGAT GTTGTACTAT 300  
GACCTGATT TGGATGATTA CCAAAATGCT CCTGGTGTCC AAACCCGGGT 350  
TCCTCATTTC GGTTGACCA AATCACTTCT ATACCTCGAC CCTCGTCTCC 400  
GGTTAGTACT TTCCAAGATA TATCATTTTG GGACATTTGC ATAATGAACA 450  
AAATAGACAT AAATTTGGGG GATTATTGTT ATATCAATAT CCATTTATAT 500  
GCTAGTCGGT AATGTGAGTG TTATGTTAGT ATAGTTAATG TGAGTGTTAT 550  
GTGATTTTCC ATTTTAAATG AAGCTAGAAA GTTGTCGTTT AATAATGTTG 600  
CTATGTCATG AGAATTATAA GGACACTATG TAAATGTAGC TTAATAATAA 650  
GGTTTGATTT GCAGAGATGC CACATCTTAC ATGGAACATT TGGTGAAAGC 700  
TCTAGAGAAA AAATGCGGGT ATGTTAACGA CCAAACCATC CTAGGAGCTC 750  
CATATGATTT CAGGTACGGC CTGGCTGCTT CGGGCCACCC GTCCCGTGTA 800  
GCCTCACAGT TCCTACAAGA CCTCAAACAA TTGGTGGAAA AAAC TAGCAG 850  
CGAGAACGAA GGAAAGCCAG TGATACTCCT CTCCCATAGC CTAGGAGGAC 900  
TTTTCGTCCT CCATTTCCTC AACCGTACCA CCCCTTCATG GCGCCGCAAG 950  
TACATCAAAC ACTTTGTTGC ACTCGCTGCG CCATGGGGTG GGACGATCTC 1000  
TCAGATGAAG ACATTTGCTT CTGGCAACAC ACTCGGTGTC CCTTTAGTTA 1050  
ACCCTTTGCT GGTCAGACGG CATCAGAGGA CCTCCGAGAG TAACCAATGG 1100  
CTACTTCCAT CTACCAAAGT GTTTCACGAC AGAACTAAAC CGCTTGTCGT 1150  
AACTCCCCAG GTTAACTACA CAGCTTACGA GATGGATCGG TTTTTTGCAG 1200  
ACATTGGATT CTCACAAGGA GTTGTGCCTT ACAAGACAAG AGTGTTCCT 1250  
TTAACAGAGG AGCTGATGAC TCCGGGAGTG CCAGTCACTT GCATATATGG 1300  
GAGAGGAGTT GATACACCGG AGGTTTTGAT GTATGGAAAA GGAGGATTCC 1350  
ATAAGCAACC AGAGATTAAG TATGGAGATG GAGATGGGAC GGTTAATTTG 1400  
GCGAGCTTAG CAGCTTTGAA AGTCGATAGC TTGAACACCG TAGAGATTGA 1450  
TGGAGTTTCC CATACTCTA TACTTAAAGA CGAGATCGCA CTTAAAGAGA 1500  
TTATGAAGCA GATTTCAATT ATTAATTATG AATTAGCCAA TGTTAATGCC 1550  
GTCAATGAAT GA 1562

<210> 11  
 <211> 3896  
 <212> genomic DNA  
 <213> Arabidopsis thaliana

<400> 11  
 ATGGGAGCGA ATTCGAAATC AGTAACGGCT TCCTTCACCG TCATCGCCGT 50  
 TTTTCTCTTG ATTTGCGGTG GCCGAAGTGC GGTGGAGGAT GAGACCGAGT 100  
 TTCACGGCGA CTACTCGAAG CTATCGGGTA TAATCATTCC GGGATTGCG 150  
 TCGACGCAGC TACGAGCGTG GTCGATCCTT GACTGTCCAT AACTCCGTT 200  
 GGAATTCAAT CCGCTCGACC TCGTATGGCT AGACACCACT AAGGTCCGTG 250  
 ATCTTCATTT CCTTCGCTCC TTATTCTGTC GGTCGAGTCA CTTGTTGATG 300  
 AATTCCAAGC GAAATATAGC AATGAAGCAT GTCTCGTCTC TCTTATTGAT 350  
 TCGTTCATTA GTCAACAGTG ACGCTTCTGA ATCTGAGTTT AGAGTCATAT 400  
 AAAACAGCTG ACTCGGCGAG TGTTCCTCCAT CGCTTTTGGT TCGCTAAATG 450  
 TAGCGCAATG AATGTGTAAT TAGTCTGCGC TTTTATTCA ACTAGATCTG 500  
 CAAGTTTTTC AGAGTGCTCA ATAGTAGTTA GAAATGTTA GGTCATTTTA 550  
 CTTGTGCATT GTGATTCTTT TGGTTGTTGC TTAGTGATCG ACGTGATGGA 600  
 TGGTTTACAG CTTCTTTCTG CTGTCAACTG CTGGTTTAAG TGTATGGTGC 650  
 TAGATCCTTA TAATCAAACA GACCATCCCC AGTGTAAGTC ACGGCCTGAC 700  
 AGTGGTCTTT CAGCCATCAC AGAATTGGAT CCAGGTTACA TAACAGGTAG 750  
 TTTCCGATTT TTCTTTCTTT TGAGTTTCT TCAATTTGAT ATCATCTTGT 800  
 TGTGATATA TATGGCTAAG TTCATTAATT TGGTCAATTT TCAGGTCTCT 850  
 TTTCTACTGT CTGGAAGAG TGGCTTAAGT GGTGTGTTGA GTTTGGTATA 900  
 GAAGCAAATG CAATTGTCGC GTTCCATAC GATTGGAGAT TGTCACCAAC 950  
 CAAATTGGAA GAGCGTGACC TTTACTTTCA CAAGCTCAAG TTAGTCCTTA 1000  
 TCAGGCTAAT GTCTTTTATC TTCTCTTTTT ATGTAAGATA AGCTAAGAGC 1050  
 TCTGGTCGTC TTCCTTTTTG CAGGTTGACC TTTGAAACTG CTTTAAACT 1100  
 CCGTGGCGGC CCTTCTATAG TATTTGCCCA TTCAATGGGT AATAATGTCT 1150  
 TCAGATACTT TCTGGAATGG CTGAGGCTAG AAATTGCACC AAAACATTAT 1200  
 TTGAAGTGCC TTGATCAGCA TATCCATGCT TATTCGCTG TTGGTACCGG 1250  
 CCTACTATCC TTAAGTTACC ATTTTATTTT TTCTCTAATT GGGGGAGTTA 1300  
 TGTGTGATCT TACTGGATTG AGCTCGATAC CTGATTTGTT GTTGATTTAG 1350  
 GAGCTCCTCT TCTTGGTTCT GTTGAGGCAA TCAAATCTAC TCTCTCTGGT 1400  
 GTAACGTTTG GCCTTCCTGT TTCTGAGGTG ACCTCTGACT TCTCTTTAGT 1450  
 TTTAAGTAGT TGATATCAAC CAGGTCTTAT AACTCACTGG ATTTTCCTTT 1500  
 TGAAAGTATT ACTTTTGTTA ATTGAAGTGC TGTACGCGAT ATGGTATCTG 1550  
 TAGATCTTGA AGTGCTAGTT ATCAAAGAAC ATATTGTGGG TAGTATACCT 1600  
 GTCAGCGGCC TTAGCTAATA CAACCAAACC ACATGTACAC TGATTTAGTT 1650  
 TTCAGATTAT TATGGTAGAC TTTAAGTTGA GAAGAACTT TGAAGTAAAT 1700  
 CTTTTTATTT TAATAGGCTA TGATTTGTTT ATTGAAATCA TGTGACATAT 1750  
 TGACATGCGC TTCTCATGTT TTTTGTGTC AAGGCTTCAG GGAAGTCTC 1800  
 GGTGTGTTGC CAATTCTTTT GCGTCGTCAT TGTGGCTTAT GCCATTTTCA 1850  
 AAGAATTGCA AGGGTGATAA CACATTCTGG ACGCATTTT CTGGGGGTGC 1900  
 TGCAAAGAAA GATAAGCGCG TATACCACTG TGATGAAGAG GAATATCAAT 1950  
 CAAAATATTC TGGCTGGCCG ACAAATATTA TTAACATTGA AATTCCTTCC 2000  
 ACTAGCGGTT AGACTCTGTA TATGCAACTG TAACACTAAC AAAAGTTTCA 2050  
 CCAAGAATGT TCACTCTCAT ATTTCTGTTCC TTTGATGTGT ATCCATCAGT 2100  
 TACAGAAACA GCTCTAGTCA ACATGACCAG CATGGAATGT GGCTTCCCA 2150  
 CCCTTTTGTC TTTACAGCC CGTGAAGTAC CAGATGGGAC TCTTTTCAA 2200  
 GCAATAGAA AGATGACCC AGATAGCAAG AGGATGTTAC ACCAGTTAAA 2250  
 GAAGTACGTA CCTTTCTTTG TGATAAGAAA TATTGCTCAT CGATCATCAC 2300  
 TTGCTGGCTT CTTGTACGTC AAATTGTTTT GTTTAAATCT CTATATCAAT 2350  
 TGTTTCATAT CTTTGTCTTT CTTACTATAA GAAACAAGTA TAATCAGAAA 2400  
 CCTTATTATT GATTATCAGT TCTCTCCTTA TATTATGGAA TGTCTTTTTC 2450  
 GTTTACAGTT ATGAATGCAA AAGGGGGTAT TTTAGTTGAT TGATCTCTC 2500  
 ATTCTCTAGT TTGTTTTGAC TAATAGCGTC AATTTTGTGT TTCTAGCAAA 2550  
 TCTTTGTGAA TTATATATAA CATGCTAACT ATACTTTTCA GGTTGTATCA 2600  
 TGATGACCCT GTTTTAAATC CTCTGACTCC TTGGGAGAGA CCACCTATAA 2650  
 AAAATGTATT TTGCATATAT GGTGCTCATC TAAAGACAGA GGTATGATGC 2700  
 ATTCTCAATA TCACATTATG CGTTGACTTT GTTATTATAT TCCCCATTTG 2750  
 GTTTGCAATA TCTTTTTGAA TTATGATTTA TCTTCTCCCT TGCATCTTAT 2800  
 GCTATTAAGC GTTAAAGGTA CTAAATGTAT GAAGCTGTCT GTCATAGGTT 2850

GGTTATTACT	TTGCCCCAAG	TGGCAAACCT	TATCCTGATA	ATTGGATCAT	2900
CACGGATATC	ATTTATGAAA	CTGAAGGTTC	CCTCGTGTCA	AGGTAATTTT	2950
CCGCAATGGC	AGAAGTAAAA	CAGGAAGGCA	AAGTCTTCTG	TATCAGTCTA	3000
GTGGCATGTT	ATCTCAGTTG	CATAAGCAAA	TTATTAAACA	ACTAAAATTT	3050
AAGTACTTTT	TTATCATTTCC	TTTTGAGCTT	AGTGGATGAT	CAGTGGCTTA	3100
AAGTGGGAAG	AGGTGTTGCA	TGAAACATGA	CAC TTGTATC	AAAGATAACT	3150
AGCAAAACAA	AACTAACCCA	TTTCTGAATT	TCATATTATT	AGGAGTAGTC	3200
GTGCTTTTAA	AAAATTTGTT	TTAAGAAACC	GAAAAACTAG	TTCATATCTT	3250
GATTGTGCAA	TATCTGCAGG	TCTGGAACCTG	TGGTTGATGG	GAACGCTGGA	3300
CCTATAACTG	GGGATGAGAC	GGTAAGCTCA	GAAGTTGGTT	TTGAAATTAT	3350
CTTCTTGCAA	ACTACTGAAG	ACTAAGATAA	TACTTGCTTC	TGGAACACTG	3400
CTTGCTATGT	TCTCTAGTAC	ACTGCAATAT	TGACTCTCCG	CTACTTTTAT	3450
TGATTATGAA	ATTGATCTCT	TATAGGTACC	CTATCATTCA	CTCTCTTGGT	3500
GCAAGAATTG	GCTCGGACCT	AAAGTTAACA	TAACAATGGC	TCCCCAGGTA	3550
CTCTTTTTTA	GTTCCCTCACC	TTATATAGAT	CAAAC TTAA	GTGTACTTTT	3600
CTGGTTATGT	GTTGATTTAC	CTCCAATTTG	TTCTTTCTAA	AAATCATATA	3650
TCTCTGTACT	CCTCAAGAAC	TTGTATTAAT	CTAAACGAGA	TTCTCATTGG	3700
GAAAATAAAA	CAACAGCCAG	AACACGATGG	AAGCGACGTA	CATGTGGAAC	3750
TAAATGTTGA	TCATGAGCAT	GGGTCAGACA	TCATAGCTAA	CATGACAAAA	3800
GCACCAAGGG	TTAAGTACAT	AACCTTTTAT	GAAGACTCTG	AGAGCATTCC	3850
GGGGAAGAGA	ACCGCAGTCT	GGGAGCTTGA	TAAAAGTGCG	TATTAA	3896

<210> 12  
<211> 709  
<212> cDNA  
<213> Lycopersicon esculentum

<400> 12  
CTGGGGCCAA AAGTGAACAT AACAAAGGACA CCACAGTCAG AGCATGATGT 50  
TCAGATGTAC AAGTGCATCT AAATATAGAG CATCAACATG GTGAAGATAT 100  
CATTCCCAAT ATGACAAAGT TACCTACAAT GAAGTACATA ACCTATTATG 150  
AGGATTCTGA AAGTTTTCCA GGGACAAGAA CAGCAGTTTG GGAGCTTGAT 200  
AAAGCAAATC ACAGGAACAT TGTCAGATCT CCAGCTTTGA TCGGGGAGCT 250  
GTGGCTTGAG ATGTGGCATG ATATTCATCC TGATAAAAAG TCCAAGTTTG 300  
TTACAAAAGG TGGTGTCTGA TCCTCACTAT TTTCTTCTAT AAATGTTTGA 350  
GTTTGTATTG ACATTGTAAG TATTGCAACA AAAAGCAAAG CGTGGGCCTC 400  
TGAGGGATGA GGA CTGCTAT TGGGATTACG GGAAAGCTCG ATGTGCATGG 450  
GCTGAACATT GTGAATACAG GTTAGAATAT TCAAATTATA TTTTGCAAAA 500  
TATTCTCTTT TTGTGTATTT AGGCCACCTT TCCCCGGTCA CAACGATGCA 550  
GATATGTATT CGGGGATGTT CACCTGGGAC AGAGTTGCAG ATTGAAGAGT 600  
TCTACATCTC ACATCCTGTC ACACTATGTG TGATATTTAA GAAACTTTGT 650  
TTGGCGGAAC AACAAAGTTTG CACAAACATT TGAAGAAGAA AGCGAAATGA 700  
TTCAGAGAG 709



&lt;210&gt; 13

&lt;211&gt; 623

&lt;212&gt; PRT

<213> *Schizosaccharomyces pombe*

&lt;400&gt; 13

MASSKKSKTHKKKKEVKSPIDLPSKKPTRLSEQPSASETQSVSNKSRKSKFGKRLNFILGAILGICGA70  
FFFAVGDDNAVFDPATLDKFGNMLGSSDLFDDIKGYLSYNVFKDAPFTTDKPSQSPSGNEVQVGLDMYNE140  
GYRSDHPVIMVPGVISSGLESWSFNCSIPYFRKRLWGSWSMLKAMFLDKQCWLEHMLDKKTGLDPKGI210  
KLRAAQGFEEADFFITGYWIWSKVIENTLAAIGYEPNNMLSASYDWRLSYANLEERDKYFSKLKMFIEYSN280  
IVHKKKVVLISHSMGSQVTTYFFKWVEAEGYGNGGPTWVNDHIEAFINISGSLIGAPKTVAALLSGEMKD350  
TGIVITLNIIEKFFSRSERAMMVRTMGGVSSMLPKGGDVAPDDLNQTNFSNGAIIRYREDIDKDHDDEFDI420  
DDALQFLKNVTDDDFKVMLAKNYSGLAWTEKEVLKNNEMPSKWINPLETSLPYAPDMKIYCVHGVGKPT490  
ERGGYYTNNPEGQPVIDSSVNDGTVENGIVMDDGDGTLPIALGLVCNKVWQTKRFNPANTSITNYEIK560  
HEPAAFDLRGGPRSAEHVDILGHSELNEIILKVSSGHGDSVPNRYISDIQEIINEINLDKPRN 623

<210> 14  
<211> 432  
<212> PRT  
<213> *Arabidopsis thaliana*

<400> 14  
MKKISSHYSVVIAILVVVTMTSMCQAVGSNVYPLILVPGNGGNQLEVRDREYKPSSVWCSSWLYPIHKK70  
SGGWFRLLWFDAAVLLSPFTRCFSDRMMLYDPPDLDDYQNAFGVQTRVPFHFGSTKSLLYLDPRLRDATSYM140  
EHLVKALEKKCGYVNDQITLGAPYDFRYGLAASGHPSRVASQFLQDLKQLVEKTSSSENEGKPVILLSHSL210  
GGLFVLHFLNRTPSWRRKYIKHFVALAAPWGGTISQMKTFAAGNTLGVPLVNPLLVRRHQRTSESNOWL280  
LPSTKVFHDRTKPLVVTPQVNYTAYEMDRFFADIGFSQGVVYPYKTRVLPLTEELMTPGVPVTCIYGRGVD350  
TPEVLMYGKGGFDKQPEIKYGDGDGTVNLAALAALKVDSLNTVEIDGVSHTSILKDEIALKEIMKQISII420  
NYELANVNAVNE 432

<210> 15  
<211> 552  
<212> PRT  
<213> *Arabidopsis thaliana*

<400> 15  
MGANSKSVTASFTVIAVFFLICGGRTAVEDETEFHGDYSKLSGIIIPGFASTQLRAWSILDCPYTFLDFN70  
PLDLVWLDTTKLLSAVNCWFKCMVLDPYNQTDHPECKSRPDSGLSAITELDPGYITGPLSTVWKEWLKWC140  
VEFGIEANAIVAVPYDWRLSPTKLEERDLYFHKLKLTFTALKLRGGPSIVFAHSMGNNVFRYFLEWLRL210  
EIAPKHLYLKWLDQHIHAYFAVGAPLLGSVEAIKSTLSGVTFGLPVSEGTARLLSNSFASSLWLMPPFSKNC280  
KGDNTFWTHFSGGAAKKDKRVYHCDEEEYQSKYSGWPTNIINIEIPSTSARELADGTLFKAIEDYDPDSK350  
RMLHQLKKYVPPFFVIRNIAHRSSLAGFLLYHDDPVFNPLTPWERPPIKNVFCIYGAHLKTEVGYYFAPSG420  
KPYPDNWIIITDIIYETEGSLVSRSGTVVDGNAGPITGDETVPYHSLSWCKNWLGPKNITMAPQILIGKI490  
KQQPEHDGSDVHVELNVDHEHGSDIIANMTKAPRVKYITFYEDSESIPGKRTAVWELDKSGY 552

5

10

15

&lt;170&gt; PatentIn Ver. 2.0

&lt;210&gt; 1Q

&lt;211&gt; 661

20 &lt;212&gt; PRT

&lt;213&gt; Saccharomyces cerevisiae

&lt;400&gt; 1

25 Met Gly Thr Leu Phe Arg Arg Asn Val Gln Asn Gln Lys Ser Asp Ser  
1 5 10 15Asp Glu Asn Asn Lys Gly Gly Ser Val His Asn Lys Arg Glu Ser Arg  
20 25 3030 Asn His Ile His His Gln Gln Gly Leu Gly His Lys Arg Arg Arg Gly  
35 40 45Ile Ser Gly Ser Ala Lys Arg Asn Glu Arg Gly Lys Asp Phe Asp Arg  
50 55 6035 Lys Arg Asp Gly Asn Gly Arg Lys Arg Trp Arg Asp Ser Arg Arg Leu  
65 70 75 8040 Ile Phe Ile Leu Gly Ala Phe Leu Gly Val Leu Leu Pro Phe Ser Phe  
85 90 95Gly Ala Tyr His Val His Asn Ser Asp Ser Asp Leu Phe Asp Asn Phe  
100 105 11045 Val Asn Phe Asp Ser Leu Lys Val Tyr Leu Asp Asp Trp Lys Asp Val  
115 120 125Leu Pro Gln Gly Ile Ser Ser Phe Ile Asp Asp Ile Gln Ala Gly Asn  
130 135 14050 Tyr Ser Thr Ser Ser Leu Asp Asp Leu Ser Glu Asn Phe Ala Val Gly  
145 150 155 16055 Lys Gln Leu Leu Arg Asp Tyr Asn Ile Glu Ala Lys His Pro Val Val  
165 170 175Met Val Pro Gly Val Ile Ser Thr Gly Ile Glu Ser Trp Gly Val Ile  
180 185 19060 Gly Asp Asp Glu Cys Asp Ser Ser Ala His Phe Arg Lys Arg Leu Trp  
195 200 205

Gly Ser Phe Tyr Met Leu Arg Thr Met Val Met Asp Lys Val Cys Trp  
 210 215 220  
 5 Leu Lys His Val Met Leu Asp Pro Glu Thr Gly Leu Asp Pro Pro Asn  
 225 230 235 240  
 Phe Thr Leu Arg Ala Ala Gln Gly Phe Glu Ser Thr Asp Tyr Phe Ile  
 245 250 255  
 10 Ala Gly Tyr Trp Ile Trp Asn Lys Val Phe Gln Asn Leu Gly Val Ile  
 260 265 270  
 Gly Tyr Glu Pro Asn Lys Met Thr Ser Ala Ala Tyr Asp Trp Arg Leu  
 275 280 285  
 15 Ala Tyr Leu Asp Leu Glu Arg Arg Asp Arg Tyr Phe Thr Lys Leu Lys  
 290 295 300  
 Glu Gln Ile Glu Leu Phe His Gln Leu Ser Gly Glu Lys Val Cys Leu  
 305 310 315 320  
 Ile Gly His Ser Met Gly Ser Gln Ile Ile Phe Tyr Phe Met Lys Trp  
 325 330 335  
 25 Val Glu Ala Glu Gly Pro Leu Tyr Gly Asn Gly Gly Arg Gly Trp Val  
 340 345 350  
 Asn Glu His Ile Asp Ser Phe Ile Asn Ala Ala Gly Thr Leu Leu Gly  
 355 360 365  
 30 Ala Pro Lys Ala Val Pro Ala Leu Ile Ser Gly Glu Met Lys Asp Thr  
 370 375 380  
 Ile Gln Leu Asn Thr Leu Ala Met Tyr Gly Leu Glu Lys Phe Phe Ser  
 385 390 395 400  
 Arg Ile Glu Arg Val Lys Met Leu Gln Thr Trp Gly Gly Ile Pro Ser  
 405 410 415  
 40 Met Leu Pro Lys Gly Glu Glu Val Ile Trp Gly Asp Met Lys Ser Ser  
 420 425 430  
 Ser Glu Asp Ala Leu Asn Asn Asn Thr Asp Thr Tyr Gly Asn Phe Ile  
 435 440 445  
 45 Arg Phe Glu Arg Asn Thr Ser Asp Ala Phe Asn Lys Asn Leu Thr Met  
 450 455 460  
 Lys Asp Ala Ile Asn Met Thr Leu Ser Ile Ser Pro Glu Trp Leu Gln  
 465 470 475 480  
 Arg Arg Val His Glu Gln Tyr Ser Phe Gly Tyr Ser Lys Asn Glu Glu  
 485 490 495  
 55 Glu Leu Arg Lys Asn Glu Leu His His Lys His Trp Ser Asn Pro Met  
 500 505 510  
 Glu Val Pro Leu Pro Glu Ala Pro His Met Lys Ile Tyr Cys Ile Tyr  
 515 520 525  
 60

Gly Val Asn Asn Pro Thr Glu Arg Ala Tyr Val Tyr Lys Glu Glu Asp  
 530 535 540  
 5 Asp Ser Ser Ala Leu Asn Leu Thr Ile Asp Tyr Glu Ser Lys Gln Pro  
 545 550 555 560  
 Val Phe Leu Thr Glu Gly Asp Gly Thr Val Pro Leu Val Ala His Ser  
 565 570 575  
 10 Met Cys His Lys Trp Ala Gln Gly Ala Ser Pro Tyr Asn Pro Ala Gly  
 580 585 590  
 Ile Asn Val Thr Ile Val Glu Met Lys His Gln Pro Asp Arg Phe Asp  
 595 600 605  
 15 Ile Arg Gly Gly Ala Lys Ser Ala Glu His Val Asp Ile Leu Gly Ser  
 610 615 620  
 20 Ala Glu Leu Asn Asp Tyr Ile Leu Lys Ile Ala Ser Gly Asn Gly Asp  
 625 630 635 640  
 Leu Val Glu Pro Arg Gln Leu Ser Asn Leu Ser Gln Trp Val Ser Gln  
 645 650 655  
 25 Met Pro Phe Pro Met  
 660  
 30 <210> 2Q  
 <211> 387  
 <212> PRT  
 <213> Arabidopsis thaliana  
 <400> 2  
 35 Val Gly Ser Asn Val Tyr Pro Leu Ile Leu Val Pro Gly Asn Gly Gly  
 1 5 10 15  
 Asn Gln Leu Glu Val Arg Leu Asp Arg Glu Tyr Lys Pro Ser Ser Val  
 20 25 30  
 40 Trp Cys Ser Ser Trp Leu Tyr Pro Ile His Lys Lys Ser Gly Gly Trp  
 35 40 45  
 45 Phe Arg Leu Trp Phe Asp Ala Ala Val Leu Leu Ser Pro Phe Thr Arg  
 50 55 60  
 Cys Phe Ser Asp Arg Met Met Leu Tyr Tyr Asp Pro Asp Leu Asp Asp  
 65 70 75 80  
 50 Tyr Gln Asn Ala Pro Gly Val Gln Thr Arg Val Pro His Phe Gly Ser  
 85 90 95  
 Thr Lys Ser Leu Leu Tyr Leu Asp Pro Arg Leu Arg Asp Ala Thr Ser  
 100 105 110  
 55 Tyr Met Glu His Leu Val Lys Ala Leu Glu Lys Lys Cys Gly Tyr Val  
 115 120 125  
 60 Asn Asp Gln Thr Ile Leu Gly Ala Pro Tyr Asp Phe Arg Tyr Gly Leu  
 130 135 140

Ala Ala Ser Gly His Pro Ser Arg Val Ala Ser Gln Phe Leu Gln Asp  
 145 150 155 160  
 5 Leu Lys Gln Leu Val Glu Lys Thr Ser Ser Glu Asn Glu Gly Lys Pro  
 165 170 175  
 Val Ile Leu Leu Ser His Ser Leu Gly Gly Leu Phe Val Leu His Phe  
 180 185 190  
 10 Leu Asn Arg Thr Thr Pro Ser Trp Arg Arg Lys Tyr Ile Lys His Phe  
 195 200 205  
 Val Ala Leu Ala Ala Pro Trp Gly Gly Thr Ile Ser Gln Met Lys Thr  
 210 215 220  
 15 Phe Ala Ser Gly Asn Thr Leu Gly Val Pro Leu Val Asn Pro Leu Leu  
 225 230 235 240  
 Val Arg Arg His Gln Arg Thr Ser Glu Ser Asn Gln Trp Leu Leu Pro  
 245 250 255  
 20 Ser Thr Lys Val Phe His Asp Arg Thr Lys Pro Leu Val Val Thr Pro  
 260 265 270  
 25 Gln Val Asn Tyr Thr Ala Tyr Glu Met Asp Arg Phe Phe Ala Asp Ile  
 275 280 285  
 Gly Phe Ser Gln Gly Val Val Pro Tyr Lys Thr Arg Val Leu Pro Leu  
 290 295 300  
 30 Thr Glu Glu Leu Met Thr Pro Gly Val Pro Val Thr Cys Ile Tyr Gly  
 305 310 315 320  
 Arg Gly Val Asp Thr Pro Glu Val Leu Met Tyr Gly Lys Gly Gly Phe  
 325 330 335  
 35 Asp Lys Gln Pro Glu Ile Lys Tyr Gly Asp Gly Asp Gly Thr Val Asn  
 340 345 350  
 40 Leu Ala Ser Leu Ala Ala Leu Lys Val Asp Ser Leu Asn Thr Val Glu  
 355 360 365  
 Ile Asp Gly Val Ser His Thr Ser Ile Leu Lys Asp Glu Ile Ala Leu  
 370 375 380  
 45 Lys Glu Ile  
 385  
 50 <210> 30  
 <211> 389  
 <212> PRT  
 <213> Arabidopsis thaliana  
 55 <400> 3  
 Leu Lys Lys Gln Gly Leu Lys Ala Lys His Pro Val Val Phe Ile Pro  
 1 5 10 15  
 60 Gly Ile Val Thr Gly Gly Leu Glu Leu Trp Glu Gly Lys Gln Cys Ala  
 20 25 30

Asp Gly Leu Phe Arg Lys Arg Leu Trp Gly Gly Thr Phe Leu Cys Trp  
 35 40 45  
 5 Val Glu His Met Ser Leu Asp Asn Glu Thr Gly Leu Asp Pro Ala Gly  
 50 55 60  
 Ile Arg Val Arg Ala Val Ser Gly Leu Val Ala Ala Asp Tyr Phe Ala  
 65 70 75 80  
 10 Pro Gly Tyr Phe Val Trp Ala Val Leu Ile Ala Asn Leu Ala His Ile  
 85 90 95  
 Gly Tyr Glu Glu Lys Asn Met Tyr Met Ala Ala Tyr Asp Trp Arg Leu  
 100 105 110  
 15 Ser Phe Gln Asn Thr Glu Arg Asp Gln Thr Leu Ser Arg Met Lys Ser  
 115 120 125  
 20 Asn Ile Glu Leu Met Val Ser Thr Asn Gly Gly Lys Lys Ala Val Ile  
 130 135 140  
 Val Pro His Ser Met Gly Val Leu Tyr Phe Leu His Phe Met Lys Trp  
 145 150 155 160  
 25 Val Glu Ala Pro Ala Pro Leu Gly Gly Gly Gly Gly Pro Asp Trp Cys  
 165 170 175  
 Ala Lys Tyr Ile Lys Ala Val Met Asn Ile Gly Gly Pro Phe Leu Gly  
 180 185 190  
 30 Val Pro Lys Ala Val Ala Gly Leu Phe Ser Ala Glu Ala Lys Asp Met  
 195 200 205  
 Arg Met Thr Arg Thr Trp Asp Ser Thr Met Ser Met Leu Pro Lys Gly  
 210 215 220  
 Gly Asp Thr Ile Trp Gly Gly Leu Asp Trp Ser Pro Glu Leu Pro Asn  
 225 230 235 240  
 40 Ala Pro Glu Met Glu Ile Tyr Ser Leu Tyr Gly Val Gly Ile Pro Thr  
 245 250 255  
 Glu Arg Ala Tyr Val Tyr Lys Leu Asn Gln Ser Pro Asp Ser Cys Ile  
 260 265 270  
 45 Pro Phe Gln Ile Phe Thr Ser Ala His Glu Glu Asp Glu Asp Ser Cys  
 275 280 285  
 Leu Lys Ala Gly Val Tyr Asn Val Asp Gly Asp Glu Thr Val Pro Val  
 290 295 300  
 Leu Ser Ala Gly Tyr Met Cys Ala Lys Ala Trp Arg Gly Lys Thr Arg  
 305 310 315 320  
 55 Phe Asn Pro Ser Gly Ile Lys Thr Tyr Ile Arg Glu Tyr Asn His Ser  
 325 330 335  
 Pro Pro Ala Asn Leu Leu Glu Gly Arg Gly Thr Gln Ser Gly Ala His  
 340 345 350  
 60 Val Asp Ile Met Gly Asn Phe Ala Leu Ile Glu Asp Ile Met Arg Val



355 360 365

Ala Ala Gly Gly Asn Gly Ser Asp Ile Gly His Asp Gln Val His Ser  
370 375 380

5 Gly Ile Phe Glu Trp  
385

10 <210> 4Q  
<211> 1986  
<212> DNA  
<213> *Saccharomyces cerevisiae*

15 <220>  
<221> CDS  
<222> (1)..(1983)

20 <400> 4  
atg ggc aca ctg ttt cga aga aat gtc cag aac caa aag agt gat tct 48  
Met Gly Thr Leu Phe Arg Arg Asn Val Gln Asn Gln Lys Ser Asp Ser  
1 5 10 15

25 gat gaa aac aat aaa ggg ggt tct gtt cat aac aag cga gag agc aga 96  
Asp Glu Asn Asn Lys Gly Gly Ser Val His Asn Lys Arg Glu Ser Arg  
20 25 30

30 aac cac att cat cat caa cag gga tta ggc cat aag aga aga agc ggt 144  
Asn His Ile His His Gln Gln Gly Leu Gly His Lys Arg Arg Arg Gly  
35 40 45

35 att agt ggc agt gca aaa aga aat gag cgt ggc aaa gat ttc gac agg 192  
Ile Ser Gly Ser Ala Lys Arg Asn Glu Arg Gly Lys Asp Phe Asp Arg  
50 55 60

40 aaa aga gac ggg aac ggt aga aaa cgt tgg aga gat tcc aga aga ctg 240  
Lys Arg Asp Gly Asn Gly Arg Lys Arg Trp Arg Asp Ser Arg Arg Leu  
65 70 75 80

45 att ttc att ctt ggt gca ttc tta ggt gta ctt ttg ccg ttt agc ttt 288  
Ile Phe Ile Leu Gly Ala Phe Leu Gly Val Leu Leu Pro Phe Ser Phe  
85 90 95

50 ggc gct tat cat gtt cat aat agc gat agc gac ttg ttt gac aac ttt 336  
Gly Ala Tyr His Val His Asn Ser Asp Ser Asp Leu Phe Asp Asn Phe  
100 105 110

55 gta aat ttt gat tca ctt aaa gtg tat ttg gat gat tgg aaa gat gtt 384  
Val Asn Phe Asp Ser Leu Lys Val Tyr Leu Asp Asp Trp Lys Asp Val  
115 120 125

60 ctc cca caa ggt ata agt tgg ttt att gat gat att cag gct ggt aac 432  
Leu Pro Gln Gly Ile Ser Ser Phe Ile Asp Asp Ile Gln Ala Gly Asn  
130 135 140

65 tac tcc aca tct tct tta gat gat ctc agt gaa aat ttt gcc gtt ggt 480  
Tyr Ser Thr Ser Ser Leu Asp Asp Leu Ser Glu Asn Phe Ala Val Gly  
145 150 155 160

70 aaa caa ctc tta cgt gat tat aat acc gag gcc aaa cat cct gtt gta 528  
Lys Gln Leu Leu Arg Asp Tyr Asn Ile Glu Ala Lys His Pro Val Val

28/53

	165	170	175	
5	atg gtc cct ggt gtc att tct acg gga att gaa agc tgg gga gtt att Met Val Pro Gly Val Ile Ser Thr Gly Ile Glu Ser Trp Gly Val Ile 180 185 190			576
10	gga gac gat gag tgc gat agt tct gcg cat ttt cgt aaa cgg ctg tgg Gly Asp Asp Glu Cys Asp Ser Ser Ala His Phe Arg Lys Arg Leu Trp 195 200 205			624
15	gga agt ttt tac atg ctg aga aca atg gtt atg gat aaa gtt tgt tgg Gly Ser Phe Tyr Met Leu Arg Thr Met Val Met Asp Lys Val Cys Trp 210 215 220			672
20	ttg aaa cat gta atg tta gat cct gaa aca ggt ctg gac cca ccg aac Leu Lys His Val Met Leu Asp Pro Glu Thr Gly Leu Asp Pro Pro Asn 225 230 235			720
25	ttt acg cta cgt gca gca cag ggc ttc gaa tca act gat tat ttc atc Phe Thr Leu Arg Ala Ala Gln Gly Phe Glu Ser Thr Asp Tyr Phe Ile 245 250 255			768
30	gca ggg tat tgg att tgg aac aaa gtt ttc caa aat ctg gga gta att Ala Gly Tyr Trp Ile Trp Asn Lys Val Phe Gln Asn Leu Gly Val Ile 260 265 270			816
35	ggc tat gaa ccc aat aaa atg acg agt gct gcg tat gat tgg agg ctt Gly Tyr Glu Pro Asn Lys Met Thr Ser Ala Ala Tyr Asp Trp Arg Leu 275 280 285			864
40	gca tat tta gat cta gaa aga cgc gat agg tac ttt acg aag cta aag Ala Tyr Leu Asp Leu Glu Arg Arg Asp Arg Tyr Phe Thr Lys Leu Lys 290 295 300			912
45	gaa caa atc gaa ctg ttt cat caa ttg agt ggt gaa aaa gtt tgt tta Glu Gln Ile Glu Leu Phe His Gln Leu Ser Gly Glu Lys Val Cys Leu 305 310 315			960
50	att gga cat tct atg ggt tct cag att atc ttt tac ttt atg aaa tgg Ile Gly His Ser Met Gly Ser Gln Ile Ile Phe Tyr Phe Met Lys Trp 325 330 335			1008
55	gtc gag gct gaa ggc cct ctt tac ggt aat ggt ggt cgt ggc tgg gtt Val Glu Ala Glu Gly Pro Leu Tyr Gly Asn Gly Gly Arg Gly Trp Val 340 345 350			1056
60	aac gaa cac ata gat tca ttc att aat gca gca ggg acg ctt ctg ggc Asn Glu His Ile Asp Ser Phe Ile Asn Ala Ala Gly Thr Leu Leu Gly 355 360 365			1104
65	gct cca aag gca gtt cca gct cta att agt ggt gaa atg aaa gat acc Ala Pro Lys Ala Val Pro Ala Leu Ile Ser Gly Glu Met Lys Asp Thr 370 375 380			1152
70	att caa tta aat acg tta gct atg tat ggt ttg gaa aag ttc ttc tca Ile Gln Leu Asn Thr Leu Ala Met Tyr Gly Leu Glu Lys Phe Phe Ser 385 390 395 400			1200
75	aga att gag aga gta aaa atg tta caa acg tgg ggt ggt ata cca tca Arg Ile Glu Arg Val Lys Met Leu Gln Thr Trp Gly Gly Ile Pro Ser 405 410 415			1248

	atg cta cca aag gga gaa gag gtc att tgg ggg gat atg aag tca tct Met Leu Pro Lys Gly Glu Glu Val Ile Trp Gly Asp Met Lys Ser Ser 420 425 430	1296
5	tca gag gat gca ttg aat aac aac act gac aca tac ggc aat ttc att Ser Glu Asp Ala Leu Asn Asn Asn Thr Asp Thr Tyr Gly Asn Phe Ile 435 440 445	1344
10	cga ttt gaa agg aat acg agc gat gct ttc aac aaa aat ttg aca atg Arg Phe Glu Arg Asn Thr Ser Asp Ala Phe Asn Lys Asn Leu Thr Met 450 455 460	1392
15	aaa gac gcc att aac atg aca tta tgg ata tca cct gaa tgg ctc caa Lys Asp Ala Ile Asn Met Thr Leu Ser Ile Ser Pro Glu Trp Leu Gln 465 470 475 480	1440
20	aga aga gta cat gag cag tac tgg ttc ggc tat tcc aag aat gaa gaa Arg Arg Val His Glu Gln Tyr Ser Phe Gly Tyr Ser Lys Asn Glu Glu 485 490 495	1488
	gag tta aga aaa aat gag cta cac cac aag cac tgg tgg aat cca atg Glu Leu Arg Lys Asn Glu Leu His His Lys His Trp Ser Asn Pro Met 500 505 510	1536
25	gaa gta cca ctt cca gaa gct ccc cac atg aaa atc tat tgt ata tac Glu Val Pro Leu Pro Glu Ala Pro His Met Lys Ile Tyr Cys Ile Tyr 515 520 525	1584
30	ggg gtg aac aac cca act gaa agg gca tat gta tat aag gaa gag gat Gly Val Asn Asn Pro Thr Glu Arg Ala Tyr Val Tyr Lys Glu Glu Asp 530 535 540	1632
35	gac tcc tct gct ctg aat ttg acc atc gac tac gaa agc aag caa cct Asp Ser Ser Ala Leu Asn Leu Thr Ile Asp Tyr Glu Ser Lys Gln Pro 545 550 555 560	1680
40	gta ttc ctc acc gag ggg gac gga acc gtt cgg ctc gtg gcg cat tca Val Phe Leu Thr Glu Gly Asp Gly Thr Val Pro Leu Val Ala His Ser 565 570 575	1728
	atg tgt cac aaa tgg gcc cag ggt gct tca cgg tac aac cct gcc gga Met Cys His Lys Trp Ala Gln Gly Ala Ser Pro Tyr Asn Pro Ala Gly 580 585 590	1776
45	att aac gtt act att gtg gaa atg aaa cac cag cca gat cga ttt gat Ile Asn Val Thr Ile Val Glu Met Lys His Gln Pro Asp Arg Phe Asp 595 600 605	1824
50	ata cgt ggt gga gca aaa agc gcc gaa cac gta gac atc ctc ggc agc Ile Arg Gly Gly Ala Lys Ser Ala Glu His Val Asp Ile Leu Gly Ser 610 615 620	1872
55	ggg gag ttg aac gat tac atc ttg aaa att gca agc ggt aat ggc gat Ala Glu Leu Asn Asp Tyr Ile Leu Lys Ile Ala Ser Gly Asn Gly Asp 625 630 635 640	1920
60	ctc gcc gag cca cgc caa ttg tct aat ttg agc cag tgg gtt tct cag Leu Val Glu Pro Arg Gln Leu Ser Asn Leu Ser Gln Trp Val Ser Gln 645 650 655	1968

1986

atg ccc ttc cca atg taa  
Met Pro Phe Pro Met  
660

5  
<210> 5 a  
<211> 661  
<212> PRT  
<213> *Saccharomyces cerevisiae*

10  
<400> 5  
Met Gly Thr Leu Phe Arg Arg Asn Val Gln Asn Gln Lys Ser Asp Ser  
1 5 10 15

15 Asp Glu Asn Asn Lys Gly Gly Ser Val His Asn Lys Arg Glu Ser Arg  
20 25 30

Asn His Ile His His Gln Gln Gly Leu Gly His Lys Arg Arg Arg Gly  
35 40 45

20 Ile Ser Gly Ser Ala Lys Arg Asn Glu Arg Gly Lys Asp Phe Asp Arg  
50 55 60

Lys Arg Asp Gly Asn Gly Arg Lys Arg Trp Arg Asp Ser Arg Arg Leu  
65 70 75 80

25 Ile Phe Ile Leu Gly Ala Phe Leu Gly Val Leu Leu Pro Phe Ser Phe  
85 90 95

Gly Ala Tyr His Val His Asn Ser Asp Ser Asp Leu Phe Asp Asn Phe  
100 105 110

Val Asn Phe Asp Ser Leu Lys Val Tyr Leu Asp Asp Trp Lys Asp Val  
115 120 125

35 Leu Pro Gln Gly Ile Ser Ser Phe Ile Asp Asp Ile Gln Ala Gly Asn  
130 135 140

Tyr Ser Thr Ser Ser Leu Asp Asp Leu Ser Glu Asn Phe Ala Val Gly  
145 150 155 160

Lys Gln Leu Leu Arg Asp Tyr Asn Ile Glu Ala Lys His Pro Val Val  
165 170 175

45 Met Val Pro Gly Val Ile Ser Thr Gly Ile Glu Ser Trp Gly Val Ile  
180 185 190

Gly Asp Asp Glu Cys Asp Ser Ser Ala His Phe Arg Lys Arg Leu Trp  
195 200 205

50 Gly Ser Phe Tyr Met Leu Arg Thr Met Val Met Asp Lys Val Cys Trp  
210 215 220

Leu Lys His Val Met Leu Asp Pro Glu Thr Gly Leu Asp Pro Pro Asn  
225 230 235 240

Phe Thr Leu Arg Ala Ala Gln Gly Phe Glu Ser Thr Asp Tyr Phe Ile  
245 250 255

60 Ala Gly Tyr Trp Ile Trp Asn Lys Val Phe Gln Asn Leu Gly Val Ile  
260 265 270

	Gly	Tyr	Glu	Pro	Asn	Lys	Met	Thr	Ser	Ala	Ala	Tyr	Asp	Trp	Arg	Leu	
			275					280					285				
5	Ala	Tyr	Leu	Asp	Leu	Glu	Arg	Arg	Asp	Arg	Tyr	Phe	Thr	Lys	Leu	Lys	
			290				295					300					
	Glu	Gln	Ile	Glu	Leu	Phe	His	Gln	Leu	Ser	Gly	Glu	Lys	Val	Cys	Leu	
	305					310					315					320	
10	Ile	Gly	His	Ser	Met	Gly	Ser	Gln	Ile	Ile	Phe	Tyr	Phe	Met	Lys	Trp	
					325					330					335		
	Val	Glu	Ala	Glu	Gly	Pro	Leu	Tyr	Gly	Asn	Gly	Gly	Arg	Gly	Trp	Val	
15				340					345					350			
	Asn	Glu	His	Ile	Asp	Ser	Phe	Ile	Asn	Ala	Ala	Gly	Thr	Leu	Leu	Gly	
			355					360					365				
20	Ala	Pro	Lys	Ala	Val	Pro	Ala	Leu	Ile	Ser	Gly	Glu	Met	Lys	Asp	Thr	
			370				375					380					
	Ile	Gln	Leu	Asn	Thr	Leu	Ala	Met	Tyr	Gly	Leu	Glu	Lys	Phe	Phe	Ser	
	385					390					395					400	
25	Arg	Ile	Glu	Arg	Val	Lys	Met	Leu	Gln	Thr	Trp	Gly	Gly	Ile	Pro	Ser	
					405					410					415		
	Met	Leu	Pro	Lys	Gly	Glu	Glu	Val	Ile	Trp	Gly	Asp	Met	Lys	Ser	Ser	
30				420					425					430			
	Ser	Glu	Asp	Ala	Leu	Asn	Asn	Asn	Thr	Asp	Thr	Tyr	Gly	Asn	Phe	Ile	
			435					440					445				
35	Arg	Phe	Glu	Arg	Asn	Thr	Ser	Asp	Ala	Phe	Asn	Lys	Asn	Leu	Thr	Met	
			450				455					460					
	Lys	Asp	Ala	Ile	Asn	Met	Thr	Leu	Ser	Ile	Ser	Pro	Glu	Trp	Leu	Gln	
	465					470					475					480	
40	Arg	Arg	Val	His	Glu	Gln	Tyr	Ser	Phe	Gly	Tyr	Ser	Lys	Asn	Glu	Glu	
					485					490					495		
	Glu	Leu	Arg	Lys	Asn	Glu	Leu	His	His	Lys	His	Trp	Ser	Asn	Pro	Met	
45				500					505					510			
	Glu	Val	Pro	Leu	Pro	Glu	Ala	Pro	His	Met	Lys	Ile	Tyr	Cys	Ile	Tyr	
			515					520					525				
50	Gly	Val	Asn	Asn	Pro	Thr	Glu	Arg	Ala	Tyr	Val	Tyr	Lys	Glu	Glu	Asp	
			530				535					540					
	Asp	Ser	Ser	Ala	Leu	Asn	Leu	Thr	Ile	Asp	Tyr	Glu	Ser	Lys	Gln	Pro	
	545					550					555					560	
55	Val	Phe	Leu	Thr	Glu	Gly	Asp	Gly	Thr	Val	Pro	Leu	Val	Ala	His	Ser	
					565					570					575		
	Met	Cys	His	Lys	Trp	Ala	Gln	Gly	Ala	Ser	Pro	Tyr	Asn	Pro	Ala	Gly	
60				580					585					590			

Ile Asn Val Thr Ile Val Glu Met Lys His Gln Pro Asp Arg Phe Asp  
595 600 605

5 Ile Arg Gly Gly Ala Lys Ser Ala Glu His Val Asp Ile Leu Gly Ser  
610 615 620

Ala Glu Leu Asn Asp Tyr Ile Leu Lys Ile Ala Ser Gly Asn Gly Asp  
625 630 635 640

10 Leu Val Glu Pro Arg Gln Leu Ser Asn Leu Ser Gln Trp Val Ser Gln  
645 650 655

Met Pro Phe Pro Met  
15 660

## SEQUENCE LISTING

&lt;110&gt; Stymne Dr., Sten

&lt;120&gt;

&lt;130&gt;

&lt;140&gt;

&lt;141&gt;

&lt;150&gt;

&lt;210&gt; 1

&lt;211&gt; 1986

&lt;212&gt; genomic DNA

<213> *Saccharomyces cerevisiae*

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1) .. (1983)

&lt;400&gt; 1

atg ggc acc ctg ttt cga aga aat gtc cag aac caa aag agt gat tct 48  
 Met Gly Thr Leu Phe Arg Arg Asn Val Gln Asn Gln Lys Ser Asp Ser  
 1 5 10 15

gat gaa aac aat aaa ggg ggt tct gtt cat aac aag cga gag agc aga 96  
 Asp Glu Asn Asn Lys Gly Gly Ser Val His Asn Lys Arg Glu Ser Arg  
 20 25 30

acc cac att cat cat caa cag gga tta ggc cat aag aga aga agg ggt 144  
 Asn His Ile His His Gln Gln Gly Leu Gly His Lys Arg Arg Arg Gly  
 35 40 45

att agt ggc agt gca aaa aga aat gag cgt ggt aaa gat ttc gac agg 192  
 Ile Ser Gly Ser Ala Lys Arg Asn Glu Arg Gly Lys Asp Phe Asp Arg  
 50 55 60

aaa aga gac ggg aac ggt aga aaa cgt tgg aga gat tcc aga aga ctg 240  
 Lys Arg Asp Gly Asn Gly Arg Lys Arg Trp Arg Asp Ser Arg Arg Leu  
 65 70 75 80

att ttc att cct ggt gca ttc tta ggt gaa ttc ttg cct ttt agc ttt 288  
 Ile Phe Ile Leu Gly Ala Phe Leu Gly Val Leu Leu Pro Phe Ser Phe  
 85 90 95

ggc ggt tat cat gtt cat aat agc gat agc gac ttg ttt gac aac ttt 336  
 Gly Ala Tyr His Val His Asn Ser Asp Ser Asp Leu Phe Asp Asn Phe  
 100 105 110

gta aat ttt gat tca ctt aaa gag tat ttg gat gat tgg aaa gat gtt 384  
 Val Asn Phe Asp Ser Leu Lys Val Tyr Leu Asp Asp Trp Lys Asp Val  
 115 120 125

ctc cca caa ggt ata agt tgg ttc att gat gat att cag gct ggt aac 432  
 Leu Pro Gln Gly Ile Ser Ser Phe Ile Asp Asp Ile Gln Ala Gly Asn  
 130 135 140

tac tcc aca tct tcc tta gat gat ctc agt gaa aat ttc gcc gtt ggt 480  
 Tyr Ser Thr Ser Ser Leu Asp Asp Leu Ser Glu Asn Phe Ala Val Gly  
 145 150 155 160

aaa caa ctc tta cgt gat tat aat acc gag gcc aaa cat cct gtt gta 528  
 Lys Gln Leu Leu Arg Asp Tyr Asn Ile Glu Ala Lys His Pro Val Val  
 165 170 175

atg gct ccc ggt gtc att tcc acg gga att gaa agc tgg gga gtt att 576  
 Met Val Pro Gly Val Ile Ser Thr Gly Ile Glu Ser Trp Gly Val Ile  
 180 185 190

gga gac gat gag tga gat agt tct gag cat ttt cgt aaa cca ctg tgg 624  
 Gly Asp Asp Glu Cys Asp Ser Ser Ala His Phe Arg Lys Arg Leu Trp  
 195 200 205

gga agt ttt taa atg ctg aga aca atg gtt atg gat aaa gct tgt tgg 672  
 Gly Ser Phe Tyr Met Leu Arg Thr Met Val Met Asp Lys Val Cys Trp  
 210 215 220

ctg aaa cat gca atg tta gat cct gaa aca ggt ctg gat cca ccg aac 720  
 Leu Lys His Val Met Leu Asp Pro Glu Thr Gly Leu Asp Pro Pro Asn  
 225 230 235 240

ttt acg cta cgt gca gca cag ggc ttc gaa tca act gat tat ttc acc 768  
 Phe Thr Leu Arg Ala Ala Gln Gly Phe Glu Ser Thr Asp Tyr Phe Ile  
 245 250 255

gca ggg tat tgg att tgg aac aaa gtt ttc caa aat ctg gga gta att 816  
 Ala Gly Tyr Trp Ile Trp Asn Lys Val Phe Gln Asn Leu Gly Val Ile  
 260 265 270

ggc tat gaa ccc aat aaa atg acc agc gct gcc tat gat tgg aag ctt 864  
 Gly Tyr Glu Pro Asn Lys Met Thr Ser Ala Ala Tyr Asp Trp Arg Leu  
 275 280 285



gca tat tta gat cca gaa aga cgc gat ags tac ttt acg aag cta aag 912  
 Ala Tyr Leu Asp Leu Glu Arg Arg Asp Arg Tyr Phe Thr Lys Leu Lys  
 290 295 300

gaa caa atc gaa ctg ttt cat caa tct agt ggt gaa aaa gtt tgt tta 960  
 Glu Gln Ile Glu Leu Phe His Gln Leu Ser Gly Glu Lys Val Cys Leu  
 305 310 315 320

att gga cat tct atg ggt tct cag att atc ttt tac ttt atg aaa cgg 1008  
 Ile Gly His Ser Met Gly Ser Gln Ile Ile Phe Tyr Phe Met Lys Trp  
 325 330 335

gtc gag gct gaa ggc cct ctt tac ggt aat ggt ggt cgt ggc tgg gtt 1056  
 Val Glu Ala Glu Gly Pro Leu Tyr Gly Asn Gly Gly Arg Gly Trp Val  
 340 345 350

aac gaa cac ata gat tca ttc att aat gca gca ggg acg ctt ctg ggc 1104  
 Asn Glu His Ile Asp Ser Phe Ile Asn Ala Ala Gly Thr Leu Leu Gly  
 355 360 365

gct cca aag gca gtt cca gct cta att agt ggt gaa atg aaa gat acc 1152  
 Ala Pro Lys Ala Val Pro Ala Leu Ile Ser Gly Glu Met Lys Asp Thr  
 370 375 380

att caa tta aat acg tta gcc atg tat ggt ttc gaa aag ttc ttc tca 1200  
 Ile Gln Leu Asn Thr Leu Ala Met Tyr Gly Leu Glu Lys Phe Phe Ser  
 385 390 395 400

aga att gag agn gca aaa atg tta caa aag tgg ggt ggt ata cca tca 1248  
 Arg Ile Glu Arg Val Lys Met Leu Gln Thr Trp Gly Gly Ile Pro Ser  
 405 410 415

atg cta cca aag gga gaa gag gtc att tgg ggg gat atg aag tca tct 1296  
 Met Leu Pro Lys Gly Glu Glu Val Ile Trp Gly Asp Met Lys Ser Ser  
 420 425 430

tca gag gat gca ttc aat aac aac act gac aca tac ggc aat ttc att 1344  
 Ser Glu Asp Ala Leu Asn Asn Asn Thr Asp Thr Tyr Gly Asn Phe Ile  
 435 440 445

cga ttt gar agt aat acg agc gat gct ttc aac aac aat ttc aca atg 1392  
 Arg Phe Glu Arg Asn Thr Ser Asp Ala Phe Asn Lys Asn Leu Thr Met  
 450 455 460

aat gag gcc att aac atg acc tta ttc aac tca cct gaa tgg ctg caa 1440  
 Lys Asp Ala Ile Asn Met Thr Leu Ser Ile Ser Pro Glu Trp Leu Gln  
 465 470 475 480

aga aga gta cat gag cag tac tgg ttc ggc tat tcc aag aat gaa gaa	1480
Arg Arg Val His Glu Gln Tyr Ser Phe Gly Tyr Ser Lys Asn Glu Glu	
485 490 495	
gag tta aga aaa aat gag cta cac cac aag cac tgg tgg aat cca atg	1536
Glu Leu Arg Lys Asn Glu Leu His His Lys His Trp Ser Asn Pro Met	
500 505 510	
gaa gta cca ctt cca gaa gct ccc cac atg aaa atc tat tgt ata tac	1584
Glu Val Pro Leu Pro Glu Ala Pro His Met Lys Ile Tyr Cys Ile Tyr	
515 520 525	
ggg gtg aac aac cca act gaa agc gca tat gta tat aag gaa gag gat	1632
Gly Val Asn Asn Pro Thr Glu Arg Ala Tyr Val Tyr Lys Glu Glu Asp	
530 535 540	
gac tcc tct gct ctg aat ttg acc atc gac tac gaa agc aag caa cct	1680
Asp Ser Ser Ala Leu Asn Leu Thr Ile Asp Tyr Glu Ser Lys Gln Pro	
545 550 555 560	
gta ttc ctg acc gag ggg gac gga acc gtt cgg ctg gtc ggc cat tca	1728
Val Phe Leu Thr Glu Gly Asp Gly Thr Val Pro Leu Val Ala His Ser	
565 570 575	
atg tgt cac aaa tgg gcc cag ggt gct tca ccg tac aac cct gcc gga	1776
Met Cys His Lys Trp Ala Gln Gly Ala Ser Pro Tyr Asn Pro Ala Gly	
580 585 590	
att aac gtt act att gtc gaa atg aaa cac cag cca gat cga ttt gat	1824
Ile Asn Val Thr Ile Val Glu Met Lys His Gln Pro Asp Arg Phe Asp	
595 600 605	
ata cgt ggt gga gca aaa agc gcc gaa cac gta gac atc ctg ggc agc	1872
Ile Arg Gly Gly Ala Lys Ser Ala Glu His Val Asp Ile Leu Gly Ser	
610 615 620	
gcs gag tgg aac gat tac atc ttg aaa att gca agc ggt aat ggc gat	1920
Ala Glu Leu Asn Asp Tyr Ile Leu Lys Ile Ala Ser Gly Asn Gly Asp	
625 630 635 640	
ctc gtc gag cca cgc caa ttg tct aat ttg agc cag tgg gtt tct cag	1968
Leu Val Glu Pro Arg Gln Leu Ser Asn Leu Ser Gln Trp Val Ser Gln	
645 650 655	

1986

atg ccc ttc cca atg taa  
Met Pro Phe Pro Met  
660

&lt;210&gt; 2

&lt;211&gt; 651

&lt;212&gt; PRT

<213> *Saccharomyces cerevisiae*

&lt;400&gt; 2

```

Met Gly Thr Leu Phe Arg Arg Asn Val Gln Asn Gln Lys Ser Asp Ser
1          5          10          15
Asp Glu Asn Asn Lys Gly Gly Ser Val His Asn Lys Arg Glu Ser Arg
20          25          30
Asn His Ile His His Gln Gln Gly Leu Gly His Lys Arg Arg Arg Gly
35          40          45
Ile Ser Gly Ser Ala Lys Arg Asn Glu Arg Gly Lys Asp Phe Asp Arg
50          55          60
Lys Arg Asp Gly Asn Gly Arg Lys Arg Trp Arg Asp Ser Arg Arg Leu
65          70          75          80
Ile Phe Ile Leu Gly Ala Phe Leu Gly Val Leu Leu Pro Phe Ser Phe
85          90          95
Gly Ala Tyr His Val His Asn Ser Asp Ser Asp Leu Phe Asp Asn Phe
100          105          110
Val Asn Phe Asp Ser Leu Lys Val Tyr Leu Asp Asp Trp Lys Asp Val
115          120          125
Leu Pro Gln Gly Ile Ser Ser Phe Ile Asp Asp Ile Gln Ala Gly Asn
130          135          140
Tyr Ser Thr Ser Ser Leu Asp Asp Leu Ser Glu Asn Phe Ala Val Gly
145          150          155          160
Lys Gln Leu Leu Arg Asp Tyr Asn Ile Glu Ala Lys His Pro Val Val
165          170          175
Met Val Pro Gly Val Ile Ser Thr Gly Ile Glu Ser Trp Gly Val Ile
180          185          190
Gly Asp Asp Glu Cys Asp Ser Ser Ala His Phe Arg Lys Arg Leu Trp
195          200          205
Gly Ser Phe Tyr Met Leu Arg Thr Met Val Met Asp Lys Val Cys Trp
210          215          220
Leu Lys His Val Met Leu Asp Pro Glu Thr Gly Leu Asp Pro Pro Asn
225          230          235          240
Phe Thr Leu Arg Ala Ala Gln Gly Phe Glu Ser Thr Asp Tyr Phe Ile
245          250          255
Ala Gly Tyr Trp Ile Trp Asn Lys Val Phe Gln Asn Leu Gly Val Ile
260          265          270
Gly Tyr Glu Pro Asn Lys Met Thr Ser Ala Ala Tyr Asp Trp Arg Leu
275          280          285
Ala Tyr Leu Asp Leu Glu Arg Arg Asp Arg Tyr Phe Thr Lys Leu Lys
290          295          300
Glu Gln Ile Glu Leu Phe His Gln Leu Ser Gly Glu Lys Val Cys Leu
305          310          315          320

```

Ile Gly His Ser Met Gly Ser Gln Ile Ile Phe Tyr Phe Met Lys Trp  
 325 330 335  
 Val Glu Ala Glu Gly Pro Leu Tyr Gly Asn Gly Gly Arg Gly Trp Val  
 340 345 350  
 Asn Glu His Ile Asp Ser Phe Ile Asn Ala Ala Gly Thr Leu Leu Gly  
 355 360 365  
 Ala Pro Lys Ala Val Pro Ala Leu Ile Ser Gly Glu Met Lys Asp Thr  
 370 375 380  
 Ile Gln Leu Asn Thr Leu Ala Met Tyr Gly Leu Glu Lys Phe Phe Ser  
 385 390 395 400  
 Arg Ile Glu Arg Val Lys Met Leu Gln Thr Trp Gly Gly Ile Pro Ser  
 405 410 415  
 Met Leu Pro Lys Gly Glu Glu Val Ile Trp Gly Asp Met Lys Ser Ser  
 420 425 430  
 Ser Glu Asp Ala Leu Asn Asn Asn Thr Asp Thr Tyr Gly Asn Phe Ile  
 435 440 445  
 Arg Phe Glu Arg Asn Thr Ser Asp Ala Phe Asn Lys Asn Leu Thr Met  
 450 455 460  
 Lys Asp Ala Ile Asn Met Thr Leu Ser Ile Ser Pro Glu Trp Leu Gln  
 465 470 475 480  
 Arg Arg Val His Glu Gln Tyr Ser Phe Gly Tyr Ser Lys Asn Glu Glu  
 485 490 495  
 Glu Leu Arg Lys Asn Glu Leu His His Lys His Trp Ser Asn Pro Met  
 500 505 510  
 Glu Val Pro Leu Pro Glu Ala Pro His Met Lys Ile Tyr Cys Ile Tyr  
 Gly Val Asn Asn Pro Thr Glu Arg Ala Tyr Val Tyr Lys Glu Glu Asp  
 530 535 540  
 Asp Ser Ser Ala Leu Asn Leu Thr Ile Asp Tyr Glu Ser Lys Gln Pro  
 545 550 555 560  
 Val Phe Leu Thr Glu Gly Asp Gly Thr Val Pro Leu Val Ala His Ser  
 565 570 575  
 Met Cys His Lys Trp Ala Gln Gly Ala Ser Pro Tyr Asn Pro Ala Gly  
 580 585 590  
 Ile Asn Val Thr Ile Val Glu Met Lys His Gln Pro Asp Arg Phe Asp  
 595 600 605  
 Ile Arg Gly Gly Ala Lys Ser Ala Glu His Val Asp Ile Leu Gly Ser  
 610 615 620  
 Ala Glu Leu Asn Asp Tyr Ile Leu Lys Ile Ala Ser Gly Asn Gly Asp  
 625 630 635 640  
 Leu Val Glu Pro Arg Gln Leu Ser Asn Leu Ser Gln Trp Val Ser Gln  
 645 650 655  
 Met Pro Phe Pro Met  
 660

<210> 36  
 <211> 2312  
 <212> genomic DNA  
 <213> Schizosaccharomyces pombe  
 <400> 3

ATGGCGTCTT CCAAGAAAGAG CAAAACATCAT AAGAAAAGAA AAGAAGTCAA	50
ATCTCCTATC GACTTACCAA ATTCAAAGAA ACCAACTCGC GCTTTGAGTG	100
AGCAACCTTC AGCGTCCGAA ACACAATCTG TTTCAAATAA ATCAAGAAAA	150
TCTAAATTTC GAAAAAGATT GAATTTTATA TTGGGCGCTA TTTTGGGAAT	200
ATCGGGTGCT TTTTTTTTCG CTGTTGAGAA CGACAATGCT GTTTTCGACC	250
CTGCTACGTT AGATAAATTT GCGAATATGC TAGGCTCTTC AGACTTGTTT	300
GATCACATTA AAGGATATTT ATCTTATAAT GTGTTTAAGG ATGCACCTTT	350
TACTACGGAC AAGCCTTCGC AGTCTCCTAG CGGAAATGAA GTTCAAGTTG	400
GTCTTGATAT GTACAATGAG GGATATCGAA GTGACCATCC TGTATTATG	450
GTTCCTGGTG TTATCAGCTC AGGATTAGAA AATTGGTCTG TTAATAATTG	500
CTCGATTCCCT TACTTTAGGA AACGTCTTTG GGGTAGCTGG TCTATGCTGA	550
AGGCAATGTT CCTTGACAAAG CAATGCTGGC TTGAACATTT AATGCTTGAT	600
AAAAAAACCG GCTTGGATCC GAAGGGAATT AAGCTGCGAG CAGCTCAGGG	650
GTTTGAGCA GCTGATTTTT TTATCACGGG CTATTGGATT TGGAGTAAAG	700
TAATTGAAAA CCTTGCTGCA ATTGATTATG AGCCTAATAA CATGTTAAGT	750
GCTTCTTACG AATTGGCGGT ATCATATGCA AATTAGAGG AACGTGATAA	800
ATATTTTTC AAGTTAAAAA TGTTCATTGA GTACAGCAAC ATTGTACATA	850
AGAAAAAGGT AGTGTTGATT TCTCACTCCA TGGGTTCACT GGTTACGTAC	900
TATTTTTTTA AATGGGTTGA AATGAGGCG TACGGAAATG GTGGACCGAC	950
TTGGGTTAAT GATCATATTG AAGCATTAT AATGTGAGT CTCGATGTT	1000
GTTTGACTAC GTTTCTAAT TTTGAATAGA TATCGGGATC TTTGATTGGA	1050
CCACCCAAAA CAGTGGCAGC GCTTTTATCG GGTGAATGA AAGATACAGG	1100
TATGTAAATT ACATTAACA TGTAAATATT TAATTTTTGC TAACCGTTT	1150
AAGTCAATT GAATCAGTT TCGGTCTATG GGTAAAGCAAT AATTGTTGA	1200
GATTGTTTAC TAATTTACTG TTTAGTTTGG AAAAATTTTT TTCCCGTTCT	1250
GAGGTATATT CAAAATACA AATGTGCTCT ACTTTTTCTA ACTTTTAATA	1300
GAGAGCCATG ATGGTTCGCA CTATGGGAGG AGTTAGTTCT ATGCTTCCTA	1350
AAGGAGGCGA TGTGTATGC GGAATGCCA GTTGGGTAAG AAATATGTGC	1400

TGTTAATTTT TTATTAATAT TTAGGCTCCA GATGATCTTA ATCAACAAA	1450
TTTTTCCAAAT GGTGCAATTA TTCGATATAG AGAAGACATT GATAAGGACC	1500
ACGATGAATT TGACATAGAT GATGCATTAC AATTTTTTAAA AAATGTTACA	1550
GATGACGATT TTAAAGTCAT CCTAGCGAAA AATTATTCCC ACGGTCTTGC	1600
TTGGACTGAA AAAGAAGTGT TAAAAAATAA CGAAATGCCG TCTAATGGA	1650
TAAATCCGCT AGAAGTAAGA ACATTAAAGT TACTAATTA TACTAACCCA	1700
AATAGACTAG TCTTCCTTAT GCTCCTGATA TGAAAATTTA TTGCGTTTAC	1750
CGGGTCGGAA AACCBACTGA GAGAGGTTAT TATTATACTA ATATCCTGA	1800
GGGGCAACCT CTCATTGATT CCTCGGTTAA TGATGGAACA AAAGTTGAAA	1850
ATGTGAGAGA ATTTATGTTT CAAACATTCT ATTAAGTGT TTATTAGGCT	1900
ATTGTTATGG ATGATGGTGA TGGAACTTTA CCAATATTAG CCCTTGGTTT	1950
GGTGCCAAAT AAAGTTTGCC AAACAAAAG GTTTAATCCT GCTAATACAA	2000
GTATCACAAA TTATGAAATC AAGCATGAAC CTGCTGCCCT TGATCTGAGA	2050
GGAGGACCTC CCTCGGCAGA ACACGTCGAT ATACTTGGAC ATTCAGAGCT	2100
AATGTATGT TCATTTTACC TTACAAATTT CTATTACTAA CTCTTGAAAT	2150
AAGGAATTA TTTTAAAAGT TTCATCAGGC CATGGTGAAT CGGTACCAA	2200
CCCTTATATA TCAGATATCC AGTACGGACA TAAGTTTTGT AGATTGCAAT	2250
TACTAACTA ACCGAACAGG GAATAATAA ATGAGATAAA TCTCGATAAA	2300
CCTAGAAATT AA	2312

<210> 4  
 <211> 3685  
 <212> genomic DNA  
 <213> Arabidopsis thaliana  
 <400> 4

ATGCCCCCTTA TTCATCGGAA AAAGCCGACG GAGAAACCAT CGACCCCGCC	50
ATCTGAAGAG GTGGTGCACG ATGAGGATTC GCAAAAGAAA CCACACGAAT	100
CTTCCAAATC CCACCATAAG AATCGAAGC GAGGAGGCA GTGGTGTGC	150
ATCGATTCTT GTTGTGTGTT CATTGGGTGT GTGTGTGTAA CCTGGTGGTT	200
TCTTCTCTTC CTTTACAACG CAATGCCTGC GAGCTTCCCT CAGTATGTAA	250
CGGAGCGAAT CACGGGTCTT TTGCCTGACC CGCCCGGTGT TAAGCTCAA	300
AAAGAAGGTC TTAAGGCGAA ACATCCTGTT GTCTTCATTC CTGGGATTGT	350
CACCGGTGGG CTCGAGCTTT GGAAGGCAA ACATGCGCT GATCGTTTAT	400
TTAGAAACG TTTGTGGGT GGAACTTTGT GTGAAGTCTA CAAAGGTGA	450
GCTCAACAAT TCTCACTCTT CCTTTATATT GGGATTGGGA TTGGATCTGA	500
TGAGATCAGC CACTTGTGTC TTCTTCAACA TCACTCAAAC TTTAATTCCA	550
TGTTTGTCTG TCTTACTCTT TACTTTTTTT TTTTTTTGAT GTGAACGCT	600
ATTTTCTTAA GAGACTATTT CTGTATGTGT AAGGTAAGCG TTCCAAGGAC	650
GTAATGGCT TCGACTATTT CTGTTTGATT GTTAACTTTA GGATATAAAA	700
TAGCTGCCCTT GGAATTTCAA GTCATCTTAT TGCCAAATCT GTTGCTAGAC	750
ATGCCCTAGA GTCCGTTTAT AACAAATTAC TTCCTTTACT GTCGTTGCGT	800
GTAGTTTAG CTTTGTGTAG CGTATAATGA AGTAGTGTTT TATGTTTTGT	850
TGGGAATAGA GAAGTTCTAA CTACATCTGT GGAAGTGTG TTCAGGCTGT	900
GATAGAGGAC TGTGCTTTA TTATTCAACT ATGTATATGT GTAATTAAG	950
CTAGTTCCTT TTTGATCTTT CAGCTCAATG TGCTTTTCTC AATTTTTTTC	1000
TCAATTTCAA AGTTTCACAT CGAGTTTATT CACATGTCTT GAATTTCTGC	1050
CATCTCGTT CTGTTATCCA GCTTTGAACT CCTCCCGACC CTGCTATGGA	1100
TATATTAAA AAAAGTGTG TTGTGGGTTG CATCTTTGTT ACGATCTGCA	1150
TCTTCTCTT TCGGCTCAGT GTTCAATGTT TTGCTATGCT AGAGATGGGC	1200
AATGTTATTG TTGATGGTAA CAGTGGTATA GTTGATAGTA TCTTAAGTAA	1250
TCAATPATCT CTTTGATCA GGCCTCTATG TTGGGTGGAA CACATGTCAC	1300
TTGACAATGA AATGGGTTG GATCCAGCTG GTATTAGAGT TCGAGCTGTA	1350



TCAGGACTCG TGGCTGCTGA CTACTTTGCT CCTGGCTACT TTGTCTGGGC	1400
AGTGCTGATT GCTAACCTTG CACATATTGG ATATGAAGAG AAAAATATGT	1450
ACATGGCTGC ATATGACTGG CGGCTTTTCT TTCAGAACAC AGAGGTTCTT	1500
TTCTCATCGT TCTTTCTATT ATTCTGTTC ATGTTACGT TCTTTCTTCA	1550
TTACTTAAGG CTIAAATATG TTTCATGTTG AATTAATAGG TACGTGATCA	1600
GACTCTTAGC CGTATGAAAA GTAATATAGA GTTGATGGTT TCTACCAACG	1650
GTGGAAAAAA AGCAGTTATA GTTCCGCATT CCATGGGGGT CTTGTATTTT	1700
CTACATTTTA TGAAGTGGGT TGAGGCACCA GCTCCTCTGG GTGGCGGGGG	1750
TGGGCCAGAT TGGTGTGCAA AGTATATTAA GGCGGTGATG AACATTGGTG	1800
GACCATTTCT TGGTGTTCOA AAAGCTGTTG CAGGGCTTTT CTCTGCTGAA	1850
GCAAGGATG TTGCAGTTGC CAGGTATTGA ATATCTGCTT ATACTTTTGA	1900
TGATCACAAC CTTGGCTCTG GAACCAAG TTATTCTACT AAATATCAAT	1950
TCTAATAACA TTGCTATATT ATCGCTGCAA CTGACATTGG TTGATTATTT	2000
TTGCTGCTTA TGTAAGTGAA ACTCTCTTGA GATTAGACAA ATGATGAATT	2050
GATAATTCTT ACCGATTGCT CTGTGATGAC CAGTTTCTTA GCTTCGACGA	2100
TAACATTTGT CATACTGTCT TTTGGAGGGC ATTGAATTTT GCTATGGAAA	2150
GCGCTGGAGC TTCCATGCTT GCATTCTTTA CCAATTAGCG TTATTCTGCT	2200
TCTTTCAATT TTCTTGATA TGCATCTATG GTCTTTTATT TCTTCTTAAT	2250
TAAAGACTCG TTGGATTAGT TGCTCTATTA GTCACCTGGT TCCTTAATAT	2300
AGACTTTTAC TTTCTTCGAA AATTGCAGAG CGATTGCCCC AGGATTCTTA	2350
GACACCGATA TATTTAGACT TCAGACCTTG CAGCATGTAA TGAGAATGAC	2400
ACGCACATGG GACTCAACAA TGTCTATGTT ACCGAAGGGA GGTGACACGA	2450
TATGGGGCGG GCTTGATTGG TCACCGGAGA AAGGCCACAC CTGTTGTGGG	2500
AAAAAGCAA AGAACAACGA AACTTGTGGT GAAGCAGGTG AAAACGGAST	2550
TTCCAAGAAA AGTCTGTGTA ACTATGGAAG GATGATATCT TTGGGGAAG	2600
AAGTAGCAGA CGCTGCGCCA TCTGAGATTA ATAATATTGA TTTTCGAGTA	2650
AGGACATATA AATCATAATA AACCTTGTAC ATTTTGTGAT TGTATGATGA	2700
ATATCTGTAC ATTTTATCTG GTGAAGGCTG CTGTCAAAGG TCAGAGTATC	2750
CCAAATCACA CCTGTCTGTA CGTGTGGACA GAGTACCATG ACATGGGAAT	2800
TGCTGGGATC AAAGCTATCG CTGASTATAA GGTCTACACT GCTGGTGAAG	2850

CTATAGATCT ACTACATTAT GTTGCTCCTA AGATGATGSC GCGTGSTGCC	2900
GCTCATTTCCT CTTATGGAAAT TGCTGATGAT TTGGATGACA CCAAGTATCA	2950
AGATCCCCAA TACTGGTCAA ATCCGTTAGA GACAAAGTAA GTGATTTCTT	3000
GATTCCAACCT GTATCCTTCG TCCTGATGCA TTATCAGTCT TTTTGTTTTT	3050
GGTCTTGTTG GATATGGTTT TCAGCTCAAA GCTTACAAAG CTGTTTCTGA	3100
GCCTTTCTCA AAAAGGCTTG CTCAGTAATA TTGAGGTGCT AAAGTTGATA	3150
CAITGACTC TTGCTTATAA ATCCTCCGTT TGGTTTGTTC TGCTTTTTC	3200
GATTACCGAA TGCTCCTGAG ATGGAAATCT ACTCATTATA CGGAGTGGGG	3250
ATACCAACGG AACGAGCATA CGTATACAAG CTTAACCAGT CTCCCGACAG	3300
TTGCATCCCC TTTCAGATAT TCACTTCTGC TCACGAGGAG GACGAAGATA	3350
GCTGTCTGAA AGCAGGAGTT TACAATGTGG ATGCGGATGA AACASTACCC	3400
GTCTTAAGTG CCGGCTACAT GTGTGCAAAA GCGTGGCCTG GCARGACAA	3450
ATTCAACCCCT TCCGGAATCA AGACTTATAT AAGAGAATAC AATCACTCTC	3500
CGCCGGCTAA CCTGTTGGAA GGSCGCGGGA CGCAGAGTGG TGCCCATGTT	3550
GATATCATGG GAAACTTTGC TTTGATCGAA GATATCATGA GCGTTGCCGC	3600
CGGAGGTAAC GGGTCTGATA TAGGACATGA CCAGGTCCAC TCTGGCATAT	3650
TTGAATGGTC GGAGCGTATT GACCTGAAGC TGTGA	3625

<210> 5  
<211> 402  
<212> cDNA  
<213> Arabidopsis thaliana  
<220>  
<221> CDS  
<222> (120)..(402)  
<400> 5

AGAAACAGCTCTTTGTCTCT CTCGACTGATCTAACAATCC CTAATCTGTGTTCTAAATTC 60  
CTGGACGAGATTTGACAAAG TCCGTATAGCTTAACCTGGT TTAATTTCAAGTGACAGAT 119  
ATG CCC CTT ATT CAT CGG AAA AAG CCG ACG CAG AAA CCA TCG ACG CCG 167  
Met Pro Leu Ile His Arg Lys Lys Pro Thr Glu Lys Pro Ser Thr Pro  
CCA TCT GAA GAG GTG GTG CAC GAT GAG GAT TCG CAA AAG AAA CCA CAC 215  
Pro Ser Glu Glu Val Val His Asp Glu Asp Ser Gln Lys Lys Pro His  
GAA TCT TCC AAA TCC CAC CAT AAG NAA TCG AAC GGA GGA GGG AAG TGG 263  
Glu Ser Ser Lys Ser His His Lys ??? Ser Asn Gly Gly Gly Lys Trp  
TCG TGC ATC GAT TCT TGT TGT TGG TTC ATT GGG TGT GTG TGT GTA ACC 311  
Ser Cys Ile Asp Ser Cys Cys Trp Phe Ile Gly Cys Val Cys Val Thr  
TGG TGG TTT CTT CTC TTC CTT TAC AAC GCA ATG CCT GCG AGC TTC CCT 359  
Trp Trp Phe Leu Leu Phe Leu Tyr Asn Ala Met Pro Ala Ser Phe Pro  
CAG TAT GTA ACG GAG CCG AAT CAC GNG TCC TTT GCC TTA CCC G 402  
Gln Tyr Val Thr Glu Pro Asn His ??? Ser Phe Ala Leu Pro

<210> 6<sup>6</sup>  
 <211> 643  
 <212> cDNA  
 <213> Zea mays  
 <220>  
 <221> CDS  
 <222> (1) .. (402)  
 <400> 6

```

CGG GAG AAA ATA GCT GCT TTG AAG GGG GGT GTT TAC TTA GCC GAT GGT  48
Arg Glu Lys Ile Ala Ala Leu Lys Gly Gly Val Tyr Leu Ala Asp Gly
   1                               10                      15

GAT GAA ACT GTT CCA GTT CTT AGT GCG GGC TAC ATG TGT GCG AAA GGA  96
Asp Glu Thr Val Pro Val Leu Ser Ala Gly Tyr Met Cys Ala Lys Gly
                20                      25                      30

TGG CST GGC AAA ACT CCT TTC AGC CCT GCG GGC AGC AAG ACT TAC GTG  144
Trp Arg Gly Lys Thr Arg Phe Ser Pro Ala Gly Ser Lys Thr Tyr Val
                35                      40                      45

AGA GAA TAC AGC CAT TCG CCA CCC TCT ACT CTC CTG GAA GGC AGG GGC  192
Arg Glu Tyr Ser His Ser Pro Pro Ser Thr Leu Leu Glu Gly Arg Gly
                50                      55                      60

ACC CAG AGC GGT GCA CAT GTT GAT ATA ATG GGG AAC TTT GCT CTA ATT  240
Thr Gln Ser Gly Ala His Val Asp Ile Met Gly Asn Phe Ala Leu Ile
   65                          70                          75                      80

GAG GAC GTC ATC AGA ATA CCT GCT GCG GCA ACC GGT GAG GAA ATT GGT  288
Glu Asp Val Ile Arg Ile Ala Ala Gly Ala Thr Gly Glu Glu Ile Gly
                85                      90                      95

GGC GAT CAG GTT TAT TCA GAT ATA TTC AAG TGG TCA GAG AAA ATC AAA  336
Gly Asp Gln Val Tyr Ser Asp Ile Phe Lys Trp Ser Glu Lys Ile Lys
                100                      105                      110

TTG AAA TTG TAA CCTATGGGAA GTTAAAGAAG TGCCGACCCG TTTATTGCGTTCC  391
Leu Lys Leu
                115

AAAGTGTCTT GCCTGAGTGC AACTCTGGAT TTTCCTTAA TATTGTAAT TTTCACGC  449

TTGATTGCTC CCTTTGTCAA ATTTACATTT GACAGGACGC CAATGCGATA CGATGTTG  507

TACCGCTATT TTCAGCATTG TATATTAAAC TGTACAGGTG TAAGTTGCAT TTGCCAGC  565

TGAAATTGTG TAGTCGTTTT CTTTACGATT TAATTANCAAG TGCCCGAGCA GTGCCCCA  623

AGCNAAAAAA AAAAAAABAA  643

```

47/53

<210> 7  
 <211> 115  
 <212> PRT  
 <213> Zee mays  
 <400> 7

Arg Glu Lys Ile Ala Ala Leu Lys Gly Gly Val Tyr Leu Ala Asp Gly  
 1 5 10 15

Asp Glu Thr Val Pro Val Leu Ser Ala Gly Tyr Met Cys Ala Lys Gly  
 20 25 30

Trp Arg Gly Lys Thr Arg Phe Ser Pro Ala Gly Ser Lys Thr Tyr Val  
 35 40 45

Arg Glu Tyr Ser His Ser Pro Pro Ser Thr Leu Leu Glu Gly Arg Gly  
 50 55 60

Thr Gln Ser Gly Ala His Val Asp Ile Met Gly Asn Phe Ala Leu Ile  
 65 70 75 80

Glu Asp Val Ile Arg Ile Ala Ala Gly Ala Thr Gly Glu Glu Ile Gly  
 85 90 95

Gly Asp Gln Val Tyr Ser Asp Ile Phe Lys Trp Ser Glu Lys Ile Lys  
 100 105 110

Leu Lys Leu  
 115

<210> 8  
<211> 616  
<212> cDNA  
<213> *Neurospora crassa*  
<400> 8

GGTGGCGAAG ACCGANGGCGG AAGTTGGAGG CTAACGAGAA TGACNCTCGG	50
AGATGGATCT ACCCTCTAGA GACACGACTA CCNTTGCACC CAGCCTCAAG	100
GTNTACNGTT TTTATGGGTA GGAAGCCGAC GGAGCGAGCC TACATCTATC	150
TGGCGCCCCG. TCCCGGGACG ACAACGCATC TTTAGATGAC GATCGATACG	200
ACTTTGACTN AGGGGCCACAT TGACCACGGT GTGATTTTGG GCGAAGGCGA	250
TGGCACAGTG AACCTTATGA GTTTGGGGTA COTGTGCAAT AAGGGGTGGA	300
AAATGACAG ATACAATCCT GCGGGCTCAA AAATAACCGT GGTGAGATG	350
CCGCATGAAC CAGAACGGTT CAATCCGAGA GGAGGGCCGA ATACGGCGGA	500
CTTAATATG TAGAAAAGGT TGAAATTTAT GAAGAGTAAT TAAATACGGC	550
ACATAGCTTA CTCAATAGTA TGAATAATA AAAAAAATT TTTTCTIAA	600
AAAAAAAAAA AAAAAA	616

<210> 9  
 <211> 1562  
 <212> genomic DNA  
 <213> *Arabidopsis thaliana*  
 <400> 9

ATGAAAAAA TATCTTCACA TTATTCCGTA GTCATAGCGA TACTCGTTGT	50
GGTGACGATG ACCTCGATGT GTCAAGCTGT GGGTAGCAAC GTGTACCCTT	100
TGATTCTGGT TCCAGGAAAC GGAGGTAACC AGCTAGAGGT ACGGCTGGAC	150
AGAGAATACA AGCCAAGTAG TGTCTGGTGT AGCAGCTGGT TATATCCGAT	200
TCATAAGAAG AGTGGTGGAT GGTTTAGGCT ATGGTTCGAT GCAGCAGTGT	250
TATTGTCTCC CTTCAACCAGG TGCTTCAGCG ATCGAATGAT GTTGTAATAT	300
GACCGTGATT TGGATGATTA CCAAAATGCT CCTGGTGTCC AATCCCGGGT	350
TCCTCATTTC GGTTCGACCA AATCACTTCT ATACTCGAC CCTCGTCTCC	400
GGTTAGTACT TTCCAAGATA TATCATTTTG GGACATTTCG ATAATGAACA	450
AAATAGACAT AAATTGGGG GATTATTGTT ATATCAATAT CCATTTATAT	500
GCTAGTCGGT AATGTGAGTG TTATGTTAGT ATAGTTAATG TGAGTGTAT	550
GTGATTTTCC ATTTTAAATG AAGCTAGAAA GTTGTGTTTT AATAATGTTG	600
CTATGTCATG AQAATTATAA GGACACTATG TAAATGTAGC TTAATAATAA	650
GGTTTGATTT GCAGAGATGC CACATCTTAC ATGGAACATT TGSTGAAAGC	700
TCTAGAGAAA AAATGCGGGT ATGTTAACGA CCAAAACCATC CTAGGAGCTC	750
CATATGATTT CAGGTACGGC CTGGCTGCTT CGGGCCACCC GTCCCGTGTA	800
GCCTCACAGT TCCTACAAGA CCTCAACAA TTGGTGGAAA AACTAGCAG	850
CGAGAACGAA GGAAAGCCAG TGATACTCCT CTCCCATAGC CTAGGAGGAC	900
TTTTCGTCTT CCATTTCCTC AACCGTACCA CCCCTTCATG GCGCCGCAAG	950
TACATCAAC ACTTTGTTGC ACTCGCTGCG CCATGGGGTG GGACGATCTC	1000
TCAGATGAAG ACATTGCTT CTGGCAACAC ACTCGGTGTC CCTTTAGTTA	1050
ACCTTTTGCT GGTGAGACGG CATCAGAGGA CCTCCGAGAG TAACCAATGG	1100
CTACTTCCAT CTACCAAGT GTTTCAGGAC AGAATAAAC CGCTTGTCGT	1150
AACTCCCCAG GTTAACTACA CAGCTTACGA GATGGATCGG TTTTTTGCAG	1200
ACATTGGATT CTCACAGGA GTTGTGCTT ACAAGACAG AGTGTGCTT	1250

TTAACAGAGG AGCTGATGAC TCCGGGAGTG CCAGTCACTT GCATATATGG	1300
GAGAGGAGTT GATACACCGG AGGTTTTTGAT GTATGGAAAA GGAGGATTCG	1350
ATAAGCAACC AGAGATTAAAG TATGGAGATG GAGATGGGAC GGTTAATTTG	1400
GCGAGCTTAG CAGCTTTGAA AGTCGATAGC TTGAACACCG TAGAGATTGA	1450
TGGAGTTTCG CATACTCTA TACTTAAAGA CGAGATCGCA CTTAAAGAGA	1500
TTATGAAGCA GATTTCATT ATTAATTATG AATTAGCCAA TGTTAATGCC	1550
GTCAATGAAT GA	1552



<210> 10  
 <211> 3896  
 <212> genomic DNA  
 <213> Arabidopsis thaliana  
 <400> 10

ATGGGAGCGA	ATTCGAAATC	AGTAACGGCT	TCCTTCACCG	TCATCGCCGT	50
TTTTTCTTG	ATTTGCGGTG	GCCGAAGTGC	GGTGGAGGAT	GAGACCGAGT	100
TTCACGGCGA	CTACTCGAAG	CTATCGGCTA	TAATCATESC	CCGATTCGS	150
TCGACGCGAGC	TACGAGCGTG	GTCGATCCTT	GACTGTCCAT	ACACTCCGTT	200
CGACTTCAAT	CCGCTCGACC	TCGTATGGCT	AGACACCACT	AAGGTCCGTG	250
ATCTTCATTT	CCTTCGCTCC	TTATTCTGTC	GGTCGAGTCA	CTTGTTGATG	300
AATTCCAAGC	GAAATATAGC	AATGAAGCAT	GTCTCGTCTC	TCATTATTGAT	350
TCGTTTCAAT	GTCAACAGTG	ACGCTTCTGA	ATCTGAGTTT	AGAGTCATAT	400
AAACAGCTO	ACTCGGCGAG	TCTTTCCCAT	CGCTTTTGGT	TCGCTAAATG	450
TAGCGCAATG	AATGTGTAAT	TAGTCTGCGC	TTTTTATTCA	ACTAGATCTG	500
CAAGTTTTTC	AGAGTGCTCA	ATAGTAGTTA	GAAATGTTA	GGTCATTTTA	550
CTGTGCAAT	GTGATTCTTT	TGGTTGTTGC	TTACTGATCG	ACGTGATGGA	600
TGGTTTACAG	CTTCTTTCTG	CTGTCAACTG	CTGGTTTAA	TGTATGGTGC	650
TAGATCCTTA	TAATCAAACA	GACCATCCCG	AGTGTAAGTC	ACGGCCTGAC	700
AGTGGTCTTT	CAGCCATCAC	AGNATTGGAT	CCAGGTTACA	TAACAGGTAG	750
TTCCGGATTT	TTCTTTCTTT	TGAGTTTTCT	TCAATTTGAT	ATCATCTTGT	800
TGTGATATA	TATGGCTAAG	TTCAATTAAT	TGGTCAATTT	TCAGGTCCTC	850
TTTCTACTGT	CTGGAAGAG	TGGCTTAAGT	GGTGTGTTGA	GTTTGGTATA	900
GAACCAATG	CAATTGTGCG	TGTTCCATAC	GATTGGAGAT	TGTCACCAAC	950
CAATTGGAA	AGCGTGACC	TTTACTTTCA	CAAGCTCAAG	TIAGTCCTTA	1000
TCAGGCTAAT	GTCTTTTATC	TTCTCTTTTT	ATGTAAGATA	AGCTAAGAGC	1050
TCTGGTCGTC	TTCTTTTTTG	CAGGTTGACC	TTTGAAACTG	CTTTAAAACT	1100
CCGTGGCGGC	CCTTCTATAG	TATTTGCCCA	TTCAATGGGT	AATAATGTCT	1150
TCAGATACTT	TCTGGAATGG	CTGAGGCTAG	AAATTGCACC	AAAACATTAT	1200
TTGAAGTGCC	TTGATCAGCA	TATCCATGCT	TATTTCCGCTG	TGGTACCGG	1250
CTTACTATCC	TTAAGTTACC	ATTTTATTTT	TTCTCTAATT	GGGGGAGTTA	1300
TGTTGTGACT	TACTGGATTG	AGCTCGATAC	CTGATTTGTT	GTTGATTTAG	1350
GAGCTCCTCT	TCTTGGTTCT	GTTGAGGCAA	TCAAATCTAC	TCTCTCTGGT	1400
GTAACGTTTTG	GCCTTCCTGT	TTCTGAGGTG	ACCTCTGACT	TCTCTTTAGT	1450
TTTAAGTAGT	TGATATCAAC	CAGCTCTTAT	AACTCACTGG	ATTTTCCTTT	1500
TGAAGTATT	ACTTTTGTTA	ATTGAACCTG	TGTACGCGAT	ATGGTATCTG	1550
TAGATCTTGA	AGTGCTAGTT	ATCAAAGAAC	ATATTGTGGG	TAGTATACCT	1600
GTCAGCGGCC	TTAGCTAATA	CAACCAAAAC	ACATGTACAC	TGATTTAGTT	1650
TTCAAGATTAT	TATGGTAGAC	TTTAAGTTGA	CAAGAAACTT	TGACTGAAAT	1700
CTTTTATTTT	TAATAGGCTA	TGATTIGTTT	ATTGAATCA	TGTGACATAT	1750
TGACATGCGC	TTCTCATGTT	TTTTGTTGGC	AAGGCTTCAG	GGAACTGCTC	1800
GGTTGTTGTC	CAATTCTTTT	GCGTCGTCAT	TGTGGCTTAT	GCCATTTTCA	1850
AAGAATTGCA	AGGGTGATAA	CACATTCTGG	ACGCATTTT	CTGGGGGTGC	1900
TGCNAGAAA	GATAAGCGCG	TATACCACTG	TGATGAAGAG	GAATATCAAT	1950
CAAAATATTC	TGGCTGGCCG	ACAAATATTA	TTAACATTGA	AATTCCTTCC	200
ACTAGCGGTT	AGACTCTGTA	TATGCAACTG	TAACACTAAC	AAAAGTTTCA	2050
CCAAGAAATG	TCACTCTCAT	ATTCCTTCC	TTTGATGTGT	ATCCATCAGT	2100
TACAGAAACA	GCTCTAGTCA	ACATGACCAG	CATGGAATGT	GGCCTTCCCA	2150
CCCTTTTGTC	TTTCAAGGCC	CGTGAAGTAG	CAGATGGGAC	TCTTTTCAAA	2200
GCAATAGAAG	ACTATGACCC	AGATAGCAAG	AGGATGTTAC	ACCAGTTAA	2250
GAAGTACOTA	CCCTTCTTTG	TGATAAGAAA	TATTGCTCAT	CGATCATCAC	2300
TTGCTGGCTT	CTTGACGTC	AAATTGTTTT	GTTTAAATCT	CTATATCAAT	2350
TGTTCAATG	CTTTGTCTTT	CTTACTATAA	GAAACAAGTA	TAATCAGAAA	2400
CCTTATTATT	GATTATCAGT	TCTCTCCTTA	TATTATGGAA	TGTCTTTTTC	2450
GTTACAGTT	ATGAATGCAA	AAGGGGGTAT	TTTAGTTGAT	TGATTCTCTC	2500
ATCTCTAGT	TTGTTTTGAC	TAATAGCGTC	AATTTTGT	TTCTAGCAAA	2550
TCTTGTGAA	TTATATATAA	CATGCTAACT	ATACTTTTCA	GGTTGTATCA	2600
TGATGACCTT	GTTTTTAATC	CTCTGACTCC	TTGGGAGAGA	CCACCTATAA	2650
AAATGTATT	TTGCATATAT	GGTGCTCATC	TAAAGACAGA	GGTATGATGC	2700
ATTCTCAATA	TCACATTATG	CGTTGACTTT	GTTATTATAT	TCCCCATTTC	2750

52/53

GTITGCAATA	TCTTTTTGAA	TTATGATTTA	TCTTCTCCCT	TGCATCTTAT	2800
CCTATTAAAGC	GTAAAGGTA	CTAAATGTAT	GAAGCTGTCT	GTCATAGGTT	2850
GGTTATTACT	TTGCCCAAG	TGGCAAACCT	TATCCTGATA	ATTGGATCAT	2900
CACGGATATC	ATTTATGAAA	CTGAAGGTTT	CCTCGTGTCA	AGGTAATTTT	2950
CCGCAATGGC	ACAAGTAAAA	CAGGAAGGCA	AAGTCTTCTG	TATCAGTCTA	3000
GTGGCATGTT	ATCTCAGTTG	CATAAGCAAA	TTATTAAACA	ACTAAAATTT	3050
AAGTACTTTT	TTATCATTC	TTTTGAGCTT	AGTGDATGAT	CAGTGGCTTA	3100
AAGTGGGAAG	AGGTGTTGCA	TGAAACATGA	CACCTTGATC	AAAGATAACT	3150
ACCAAAACAA	AACTAACCCA	TTTCTGAATT	TCATATTATT	AGGAGTAGTC	3200
CTGCTTTTAA	AAAATTTGTT	TTAAGAAACC	GAAAAACTAG	TTCATATCTT	3250
GATTGTGCAA	TATCTGCAGG	TCTGGAACTG	TGGTTGATGG	GAACGCTGGA	3300
CCTATAACTG	GGGATGAGAC	GGTAAGCTCA	GAAGTTGGTT	TTGAAATTAT	3350
CTTCTTGCAA	ACTACTGAAG	ACTAAGATAA	TACTTGCTTC	TGGAACACTG	3400
CTTGCTATGT	TCTCTAGTAC	ACTGCAATAT	TGACTCTCCG	CTACTTTTAT	3450
TGATTATGAA	ATTGATCTCT	TATAGGTACC	CTATCATTCA	CTCTCTTGGT	3500
GCAAGAATTG	GCTCGGACCT	AAAGTTAACA	TAACAATGGC	TCCCCAGGTA	3550
CTCTTTTTTA	GTTCCCTCACC	TTATATAGAT	CAAACCTTAA	GTGTACTTTT	3600
CTGGTTATGT	GTTGATTTAC	CTCCAATTTG	TTCTTTCTAA	AAATCATATA	3650
TCCTGTACT	CCTCAAGAAC	TTGTATTAA	CTAAACGAGA	TTCTCATTGG	3700
GAAATATAAA	CAACAGCCAG	AACACGATCG	AAGCGACGTA	CATGTGGAAC	3750
TAAATGTTGA	TCATGAGCAT	GGGTCAGACA	TCATAGCTAA	CATGACAAAA	3800
GCAACCAAGGG	TTAAGTACAT	AACCTTTTAT	GAAGACTCTG	AGAGCATTCC	3850
GGGGAAGAGA	ACCGCAGTCT	CGGAGCTTGA	TAAAAGTGGG	TATTAA	3896

<210> 11<sup>h</sup>  
<211> 709  
<212> cDNA  
<213> tomato  
<400> 11

CTGGGGCCAA AAGTGAACAT AACAAAGGACA CCACAGTCAG AGCATGATGT	50
TCAGATGTAC AAGTGCATCT AATATAGAG CATCAACATG GTGAAGATAT	100
CATCCCAAT ATGACAAAGT TACCTACAAT GAAGTACATA ACCTATTATG	150
AGGATTCTCA AAGTTTTCCA GGGACAAGAA CAGCAGTTTG GGAGCTTGAT	200
AAAGCAAATC ACAGGAACAT TGTCAGATCT CCAGCTTTGA TGCGGGAGCT	250
GTGGCTTGAG ATGTGGCATG ATATTCATCC TGATAAAAAG TCCAAGTTTG	300
TTACAAAAGG TGGTGTCTGA TCCTCACTAT TTTCTTCTAT AATGTTTGA	350
GTGTGTATTG ACATTGTAAG TATTGCAACA AAAAGCAAAG CGTGGGCCTC	400
TCAGGGATCA GGAATGCTAT TGGGATTACG GGAAAGCTCG ATGTGCATGG	450
GCTGAACATT GTGAATACAG GTTACAATAT TCATATTATA TTTTGCAAAA	500
TATTCTCTTT TTGTGTATTT AGGCCACCTT TCCCCGGTCA CAACGATGCA	550
GATATGTATT CGGGGATGTT CACCTGGGAC AGAGTTGCAG ATTGAAGAGT	600
TCTACATCTC ACATCCTGTC AACTATGTG TGATATTTAA GAAACTTTGT	650
TTGGCGGAAC AACAAAGTTTG CACAAACATT TGAAGAAGAA AGCGAAATGA	700
TTGAGAGAG	709

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
12 October 2000 (12.10.2000)

PCT

(10) International Publication Number  
**WO 00/60095 A3**(51) International Patent Classification<sup>7</sup>: **C12N 15/54**,  
9/10, 15/81, 15/82, 1/16, 5/10, A01K 67/027, C12P 7/64(21) International Application Number: **PCT/EP00/02701**(22) International Filing Date: **28 March 2000 (28.03.2000)**(25) Filing Language: **English**(26) Publication Language: **English**(30) Priority Data:  
99106656.4 1 April 1999 (01.04.1999) EP  
99111321.8 10 June 1999 (10.06.1999) EP  
60/180,687 7 February 2000 (07.02.2000) US(71) Applicant (for all designated States except US): **BASF PLANT SCIENCE GMBH [DE/DE]; D-67056 Ludwigshafen (DE).**

(72) Inventors; and

(75) Inventors/Applicants (for US only): **DAHLQVIST, Anders [SE/SE]; Hemmansvägen 2, S-244 66 Furulund (SE). STAHL, Ulf [SE/SE]; Liljegatan 7b, S-753 24 Uppsala (SE). LENMAN, Marit [SE/SE]; Revingegatan 13a, S-223 59 Lund (SE). BANAS, Antoni [SE/PL];**Wiolinowa 14, PL-08110 Siedlce (PL). **RONNE, Hans [SE/SE]; Dirigentvägen 169, S-756 54 Uppsala (SE). STYMNE, Sten [SE/SE]; Torrlösa 1380, S-269 90 Svalöv (SE).**(74) Agent: **FITZNER, Uwe; Lintorfer Str. 10, D-40878 Ratingen (DE).**


(81) Designated States (national): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— With international search report.

[Continued on next page]

(54) Title: **ENZYMES OF THE BIOSYNTHETIC PATHWAY FOR THE PRODUCTION OF TRIACYLGLYCEROL AND RECOMBINANT DNA MOLECULES ENCODING THESE ENZYMES**

(57) Abstract: The present invention relates to the isolation, identification and characterization of nucleotide sequences encoding an enzyme catalysing the transfer of fatty acids from phospholipids to diacylglycerol in the biosynthetic pathway for the production of triacylglycerol, to the said enzymes and a process for the production of triacylglycerols.

WO 00/60095 A3



(88) Date of publication of the international search report:  
1 February 2001

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*